

From the Department of Laboratory Medicine
Karolinska Institutet, Stockholm, Sweden

Cellular mechanisms of human muscle fatigue and disease-related loss of muscle function

Karl Olsson



**Karolinska
Institutet**

Stockholm 2020

All previously published papers were reproduced with permission from the publisher.

Published by Karolinska Institutet.

Printed by Universitetsservice US-AB

© Karl Olsson, 2020

ISBN 978-91-7831-878-0

Cellular mechanisms of human muscle fatigue and disease-related loss of muscle function

THESIS FOR DOCTORAL DEGREE (Ph.D.)

By

Karl Olsson

Principal Supervisor:

Professor Thomas Gustafsson
Karolinska Institutet
Department of Laboratory Medicine
Division of Clinical Physiology

Opponent:

Professor William E. Louch
University of Oslo
Institute for Experimental Medical
Research

Co-supervisors:

Eric Rullman, MD, PhD
Karolinska Institutet
Department of Laboratory Medicine
Division of Clinical Physiology

Examination Board:

Professor Tore Bengtsson
Stockholm University
Department of Molecular Biosciences

Associate Professor Johanna Lanner
Karolinska Institutet
Department of Physiology and
Pharmacology

Associate Professor Elodie Ponsot
Örebro University
School of Health Sciences

Professor Olav Rooyackers
Karolinska Institutet
Department of CLINTEC
Division of Anaesthesiology and
Intensive Care

Associate Professor Kristian Vissing
Aarhus University
Department of Public Health - Sport
Science

Abstract

A loss of skeletal muscle function and mass is a hallmark of several diseases resulting in a dismal life quality and poor prognosis. Impaired intracellular Ca^{2+} -handling has been proposed as an underlying cause of these manifestations. This thesis investigated the cellular mechanisms of human muscle fatigue and disease-related loss of muscle function and mass. Special focus was put on the contribution of altered intracellular Ca^{2+} -handling to these processes. The thesis makes three main contributions by: (i) providing mechanistic insight into the underlying causes of fatigue in human muscle fibres, (ii) elucidating the direct effects of vitamin D in human muscle, and (iii) implicating sphingomyelinase (SMase) activation in disease-related loss of muscle function and mass.

Study I examined single intact muscle fibres and differentiated muscle stem cells (myotubes) as experimental models of human skeletal muscle. Results revealed that single intact fibres could be manually dissected, enabling studies of Ca^{2+} -regulated processes and force generation. Major differences in intracellular Ca^{2+} -handling between myotubes and muscle fibres were discerned. In **study II**, the effects of vitamin D in human skeletal muscle were investigated. Data demonstrated a modulatory role of vitamin D on muscle stem cells, but did not support presence of the vitamin D receptor in muscle fibres. In **study III**, the cellular mechanisms of fatigue were examined in single intact fibres. Fatigue-induced force loss was caused by a reduction in sarcoplasmic reticulum (SR) Ca^{2+} release, while a decrease in myofibrillar Ca^{2+} sensitivity played no role. Acidification did not reduce force production or fatigue tolerance of human fibres. **Study IV** investigated the tentative role of SMase in disease-related muscle weakness and atrophy. SMase produced a prompt force decline by a reduction in SR Ca^{2+} release and decreased myofibrillar Ca^{2+} sensitivity, while promoting atrophy processes. Intramuscular SMase activity was elevated in heart failure compared to control individuals, and positively correlated with circulating markers of inflammation and atrophy, and with factors carrying prognostic value.

This thesis reveals differences in the intracellular Ca^{2+} -handling properties of myotubes and muscle fibres, and in the underlying causes of fatigue in humans and animals. This warrants caution when transferring findings in animal fibres and human myotubes to the situation in human skeletal muscle. The effects of vitamin D on muscle stem cells suggest an importance of vitamin D in skeletal muscle health. This should be considered in conditions of severe vitamin D deficiency, e.g. chronic kidney disease. Results implicate elevated SMase activity in the advent of muscle weakness and atrophy, suggesting SMase as a tentative target for therapeutic interventions in diseases associated with a loss of skeletal muscle function and mass.

List of scientific papers

This thesis is based on the following papers, herein referred to by their Roman numerals.

- I. **Olsson K**, Cheng AJ, Alam S, Al-Ameri M, Rullman E, Westerblad H, Lanner JT, Bruton JD, Gustafsson T. Intracellular Ca^{2+} -handling differs markedly between intact human muscle fibres and myotubes. *Skeletal Muscle*. 2015;5:26. DOI 10.1186/s13395-015-0050-x
- II. **Olsson K**, Saini A, Strömberg A, Alam S, Lilja M, Rullman E, Gustafsson T. Evidence for vitamin D receptor expression and direct effects of $1\alpha,25(\text{OH})_2\text{D}_3$ in human skeletal muscle precursor cells. *Endocrinology*. 2016;157(1):98-111.
- III. **Olsson K**, Cheng AJ, Al-Ameri M, Wyckelsma VL, Rullman E, Westerblad H, Lanner JT, Gustafsson T, Bruton JD. Impaired sarcoplasmic reticulum Ca^{2+} release is the major cause of fatigue-induced force loss in intact single fibres from human intercostal muscle. *The Journal of Physiology*. 2020;598(4):773–787.
- IV. **Olsson K**, Cheng AJ, Al-Ameri M, Tardif N, Melin M, Rooyackers O, Lanner JT, Westerblad H, Gustafsson T, Bruton JD, Rullman E. Elevated sphingomyelinase activity in heart failure patients depresses skeletal muscle fibre force production and promotes atrophy. *Manuscript*.

Acknowledgements

Often a seemingly lonely (and endless) process, this thesis would likely never have been completed and, if so, with both the thesis and myself in a far worse state than the present without the support from supervisors, colleagues, friends and family. It is impossible to thank everyone involved and this acknowledgment will therefore inevitably be incomprehensive. I blame this partly on limited space and partly on my notoriously bad memory. To all those not included, I express not only my sincere gratitude but also my deepest apologies.

To my supervisors: Thomas Gustafsson and Eric Rullman, thank you for taking me on as a naïve medical student and aspiring physiologist. Thomas, Eric, Johanna Lanner and Olav Rooyackers, thanks for your including attitude and for providing a creative and fun atmosphere, both inside and outside the lab.

Arthur Cheng and Joseph Bruton, I am grateful for your enthusiasm to test the idea of dissecting single intact human fibres and all the subsequent cumbersome hours spent dissecting, both day and night. Your exceptional skills in manual dissections, force and intracellular Ca^{2+} measurements, as well as intellectual contributions have been imperative. I have thoroughly enjoyed your company and the discussions of various topics we've had during these years. Thank you Håkan Westerblad, for welcoming me into your lab and for valuable discussions. For the help with data gathering and intellectual contributions, I am grateful to my co-authors: Seher Alam, Mamdoh Al-Ameri, Victoria Wyckelsma, Mats Lilja, Amarjit Saini, Nicolas Tardif, Michael Melin and Anna Strömberg. To all members of the clinical physiology lab, thank you for creating an enjoyable and inspirational work environment.

To my parents, thank you for your unrelenting love and belief in me during my various projects, both in academia and in the mountains. Maria, thank you for your love and encouragement, despite me often making it difficult for you to do so. How you have managed to support me through this endeavour while simultaneously finalizing your own thesis and remaining a devoted mother is incomprehensible. Despite the ups and downs, we made it. I look forward to embarking on the journey towards all the adventures that still lie ahead of us. To Bertil, our son, thank you for the countless laughs, tears, sleepless nights, insights and perspectives on what truly matters in life. I hope I can be an inspiration for you to go climb your own mountain.

Karl Olsson
August 2020

Contents

Abstract.....	I
List of scientific papers	III
Acknowledgements	V
1 Introduction and background	1
2 Objectives and outline of the thesis	13
3 Summary of study results	15
Study I: <i>Ex vivo</i> models of human skeletal muscle	15
Study II: Effects of vitamin D in human skeletal muscle	16
Study III: Cellular mechanisms of human skeletal muscle fatigue.....	19
Study IV: The role of sphingomyelinase in disease-related loss of muscle function and mass.....	22
4 Study I	25
5 Study II	47
6 Study III.....	73
7 Study IV	97
8 Discussion and implications	115
<i>Ex vivo</i> models of human skeletal muscle	115
Effects of vitamin D in human skeletal muscle.....	116
Cellular mechanisms of fatigue in human muscle fibres	117
Acidosis and human muscle fatigue	118
The role of sphingomyelinase in disease-related loss of muscle function and mass	120
References	122

1 Introduction and background

Several human diseases are associated with a high incidence of a profound and debilitating loss of skeletal muscle function and mass that result in a dismal life quality and poor prognosis (Anker *et al.*, 1997; Hulsmann *et al.*, 2004; Sokka *et al.*, 2008; Chung *et al.*, 2014; Powers *et al.*, 2016). More than just the result of muscle disuse, disease-related changes inside the skeletal muscle fibres contribute to the progressive skeletal muscle dysfunction (Ward *et al.*, 2003; Powers *et al.*, 2016; Steinz *et al.*, 2019). Experimental evidence from animal models has implicated dysfunctional intracellular Ca^{2+} -handling with the impaired muscle function (Reiken *et al.*, 2003; Ward *et al.*, 2003; Yamada *et al.*, 2015b; Llano-Diez *et al.*, 2016). Factors suggested to contribute to the changes in intracellular Ca^{2+} -handling include increased neuroendocrine activation, elevated circulating levels of pro-inflammatory cytokines, and altered sphingolipid metabolism (Reiken *et al.*, 2003; Andersson & Marks, 2010; Nikolova-Karakashian & Reid, 2011; Reid & Moylan, 2011; Stasko *et al.*, 2013). Meanwhile, vitamin D deficiency is a common finding in diseased individuals associated with muscle weakness and atrophy, ostensibly via both Ca^{2+} -dependent and Ca^{2+} -independent processes (Girgis *et al.*, 2013). The current thesis investigated the intracellular mechanisms of human muscle fatigue and disease-related loss of muscle function and mass. A special focus was put on the role of altered intracellular Ca^{2+} -handling in these processes.

Skeletal muscle force development

Skeletal muscle contraction occurs when the contractile proteins myosin and actin in the force generating units, the sarcomeres, interact to form cycling cross bridges. In this reaction, ATP serves as an energy-carrying compound while being converted to ADP and free organic phosphate (P_i). In the resting muscle fibre, the myofilament regulatory protein tropomyosin forms a complex together with troponin C, T and I that blocks the myosin binding-sites on actin. A conformational change in tropomyosin, instigated by Ca^{2+} binding to troponin C, enables myosin and actin interaction. Elevations in the myoplasmic free Ca^{2+} concentration ($[\text{Ca}^{2+}]_i$) promote Ca^{2+} binding to troponin C, while Ca^{2+} dissociation from troponin C is promoted as $[\text{Ca}^{2+}]_i$ is lowered to its resting value (Gordon *et al.*, 2000). Intracellular Ca^{2+} thus serves as the cellular signalling molecule that dictates initiation and termination of muscle contraction.

The level of $[Ca^{2+}]_i$ determines the number of cross bridges that can be formed and thus regulates the amount of force development within individual skeletal muscle fibres (Gordon *et al.*, 2000). In this way, increasing levels of $[Ca^{2+}]_i$ results in larger force developments until all cross bridges are activated and the maximal Ca^{2+} -activated force is produced. There is a strong correlation between muscle fibre diameter and potential force development (Schantz *et al.*, 1983). Muscle fibre diameter is determined by the balance between protein synthesis and degradation, but may also be modulated by the activity of muscle stem cells, as will be discussed further in the following sections.

Skeletal muscle excitation-contraction coupling

The events from sarcolemmal depolarisation to initiation of cross bridge cycling are described as excitation-contraction coupling. Depolarisation of the sarcolemma is initiated at the neuromuscular junction when acetylcholine is released by the α -motoneuron (Rebbbeck *et al.*, 2014). Acetylcholine binds to receptors at the sarcolemma triggering the opening of ion channels that allow passage of both Na^+ and K^+ . This results in a localized depolarisation of the sarcolemma, that when it reaches a threshold value opens voltage-gated Na^+ channels (Nav1.4) generating an action potential that is propagated along the fibre via the sarcolemma and transverse tubular system (Gordon *et al.*, 2000). In the transverse tubules, L-type voltage-activated Ca^{2+} channels (dihydropyridine receptor; DHPR) are activated by the propagating action potential. In skeletal muscle, DHPR mechanically activates the sarcoplasmic reticulum (SR) Ca^{2+} channels (type 1 ryanodine receptor; RyR1) (Tanabe *et al.*, 1990; Rebbbeck *et al.*, 2014). At rest, the SR lumen contains 10-20 mM Ca^{2+} , the majority of which is bound to calsequestrin, while $[Ca^{2+}]_i$ is only about 50 nM. This large concentration gradient allows rapid elevation of $[Ca^{2+}]_i$ following RyR1 activation and opening (Allen *et al.*, 2008).

When α -motoneuron activation ceases the sarcolemma is repolarized, RyR1 is inactivated, and SR Ca^{2+} release stops. The re-uptake of Ca^{2+} into the SR by the SR Ca^{2+} -ATPase (SERCA) returns $[Ca^{2+}]_i$ to resting levels. Ca^{2+} dissociates from troponin C causing tropomyosin to again block the myosin binding-sites on actin whereby cross bridge cycling terminates (Gordon *et al.*, 2000).

Intracellular Ca^{2+} in skeletal muscle physiology and pathology

Intracellular Ca^{2+} -handling refers to the cellular regulation of $[\text{Ca}^{2+}]_i$ homeostasis, including the regulation of basal levels and fluctuations in $[\text{Ca}^{2+}]_i$, and the cellular response to fluctuations in $[\text{Ca}^{2+}]_i$. Intracellular Ca^{2+} -handling involves a large number of intracellular membrane or organelle bound proteins, as well as soluble cytosolic or organelle-confound proteins (Berchtold *et al.*, 2000). In skeletal muscle, intracellular Ca^{2+} -handling is a key regulator and intracellular signalling molecule implicated in muscle force generation, fatigue development, regulation of gene expression and cellular metabolism (Berchtold *et al.*, 2000; Lanner *et al.*, 2006; Gundersen, 2011; Tavi & Westerblad, 2011).

Changes in $[\text{Ca}^{2+}]_i$ transient amplitude and duration, as well as frequency of the $[\text{Ca}^{2+}]_i$ transients, modulate the intracellular response to fluctuations in $[\text{Ca}^{2+}]_i$ (Dolmetsch *et al.*, 1997; Tavi & Westerblad, 2011). Based on studies in animal and cell models, changes in baseline $[\text{Ca}^{2+}]_i$ have been implicated in both the physiological adaptation to exercise (e.g. mitochondrial biogenesis) (Bruton *et al.*, 2010; Place *et al.*, 2015) and maladaptation in disease (e.g. muscle weakness, fatigue and atrophy) (Bhattacharyya *et al.*, 1993; Menconi *et al.*, 2004; Smith & Dodd, 2007; Dridi *et al.*, 2020; Kushnir *et al.*, 2020).

Transient and prolonged changes in skeletal muscle intracellular Ca^{2+} -handling may be induced by several physiological and pathological stimuli. These include stimuli such as acute exercise and training, muscle disuse, hormonal and inflammatory factors (Berchtold *et al.*, 2000; Allen *et al.*, 2016). Transient changes (seconds to hours) in intracellular Ca^{2+} -handling are generally the result of changes in the levels of ions and metabolites that regulate the activity of Ca^{2+} -handling proteins. Prolonged changes (hours to days) may be caused by posttranslational modifications, changes in protein expression or stability of Ca^{2+} -handling proteins, or changes in the binding of regulatory proteins to Ca^{2+} -handling proteins (Berchtold *et al.*, 2000; Place *et al.*, 2015).

Skeletal muscle fatigue

Skeletal muscle fatigue has been defined as the reversible reduction in maximal voluntary force/power-producing capacity following prolonged or repeated contractions (Gandevia, 2001; Place *et al.*, 2010). A distinction between central and peripheral components limiting force production in fatigue is often made. Central fatigue refers to processes within the nervous system whereas peripheral fatigue refers to processes at, or peripheral to, the neuromuscular junction (Taylor *et al.*, 2006). It stands clear that both components participate to development of

fatigue, but the relative contribution of the two differs depending on the duration and frequency of contractions, duty cycle and external milieu (Gandevia, 2001). It is generally recognized that in the majority of situations a large part of fatigue development stems from peripheral components (Allen *et al.*, 2008).

Cellular mechanisms of skeletal muscle fatigue

Studies in isolated intact and skinned mammalian muscle fibres have demonstrated three changes to ultimately cause the fatigue-induced force loss. These are (i) a reduction in myofibrillar maximal Ca^{2+} -activated force, (ii) decreased myofibrillar Ca^{2+} sensitivity, and (iii) reduced SR Ca^{2+} release (Westerblad & Allen, 1991, 1993a; Allen *et al.*, 2008) (**Figure 1.1**). The relative importance of these changes differs during three distinctly identifiable phases during fatigue; an initial phase of depressed force while tetanic $[\text{Ca}^{2+}]_i$ increases, explained by a reduced myofibrillar maximal Ca^{2+} -activated force; a second phase where tetanic $[\text{Ca}^{2+}]_i$ and force is little affected; followed by a final phase in which both tetanic $[\text{Ca}^{2+}]_i$ and force rapidly declines. This third phase is explained by a concurrent reduction in SR Ca^{2+} release and decreased myofibrillar Ca^{2+} sensitivity (Westerblad & Allen, 1991, 1993a; Allen *et al.*, 2008).




Figure 1.1. Illustration of the cellular mechanisms of fatigue in rodent muscle fibres. A, force and $[\text{Ca}^{2+}]_i$ in the 1st, 10th and last (50th) tetanus of fatigue. Note the initial increase in tetanic $[\text{Ca}^{2+}]_i$ while force decreases between the 1st and 10th tetanus. B, proposed sequence of events during fatigue development (red line) and mechanisms causing fatigue in rodent muscle fibres: (i) a reduction in myofibrillar maximal Ca^{2+} -activated force, (ii) decreased myofibrillar Ca^{2+} sensitivity, and (iii) reduced tetanic $[\text{Ca}^{2+}]_i$ due to impaired SR Ca^{2+} release. Force- $[\text{Ca}^{2+}]_i$ relation in unfatigued (solid line) and fatigued (dashed line) fibres. Redrawn from Allen, Lamb & Westerblad; J Appl Physiol 104: 296–305, 2008.

Accumulation of P_i is believed to be of fundamental importance to the intrinsic skeletal muscle changes that ultimately causes the force depression during fatigue. The initial decline in myofibrillar maximal Ca^{2+} -activated force is not seen in mouse muscle fibres deficient of creatine kinase where there is little change in P_i (Dahlstedt *et al.*, 2000), and P_i depresses tetanic force in skinned fibres (Pate & Cooke, 1989; Millar & Homsher, 1990). The reduction in tetanic $[Ca^{2+}]_i$ is believed to mainly stem from perturbed SR Ca^{2+} release resulting from Ca^{2+} - P_i precipitation in the SR (Fryer *et al.*, 1995; Westerblad & Allen, 1996), and the combined effects of reduced cytosolic [ATP] and increased cytosolic free $[Mg^{2+}]$, which inhibits the RyR1 channel (Blazev & Lamb, 1999). Accordingly, the fall in tetanic $[Ca^{2+}]_i$ is greatly delayed when creatine kinase is inhibited in rodent fibres (Dahlstedt & Westerblad, 2001), and injection of P_i into single intact mouse fibres causes a marked fall in SR Ca^{2+} release and tetanic force (Westerblad & Allen, 1996). Reduced myofibrillar Ca^{2+} sensitivity is caused by an accumulation of P_i and a reduction in intracellular pH (pH_i), but may also be modulated by reactive oxygen and nitrogen species (RONS) (Dahlstedt *et al.*, 2001; Allen *et al.*, 2008; Lamb & Westerblad, 2011).

Intramuscular acidosis and skeletal muscle fatigue development

In exercising humans, pH_i has been reported to fall ~0.5 pH units following intense muscle activity (Sahlin *et al.*, 1975; Sahlin *et al.*, 1976; Bangsbo *et al.*, 1996). Historically, intramuscular acidosis has been acclaimed a major role in the force decline during fatigue (Fitts, 2016). The underlying observations supporting this view include observations from exercising humans and studies from isolated animal muscle fibres.

Initial studies found a good temporal correlation between intramuscular acidification and fatigue (Dawson *et al.*, 1978). In coherence with this finding, a reduction in maximal Ca^{2+} -activated force was demonstrated upon acidification of isolated animal muscles (Donaldson *et al.*, 1978; Fabiato & Fabiato, 1978). This notion was later challenged as the force-depressive effect of acidification was found to be highly temperature-dependent and to be blunted when studied at physiological temperatures (Pate *et al.*, 1995; Westerblad *et al.*, 1997). Notably, the fatigue tolerance of isolated rodent fibres fatigued under acidic conditions was unaltered when studied at near-physiological temperatures (Bruton *et al.*, 1998).

Recovery from fatigue

Recovery of force producing capacity following fatigue follows a time course that is largely dependent on the metabolic properties of the muscle fibre. While oxidative slow-twitch fibres recover their force producing capacity nearly instantly, fast-twitch fibres recover more slowly (Allen *et al.*, 2008). In fast-twitch

fibres it is possible to discern a difference in the time course for recovery depending on the frequency of stimulation used to initiate contraction. At high-frequency stimulation (80-100 Hz) recovery from fatigue occurs within minutes, while at low-frequency stimulation (20-30 Hz) recovery is not complete until several hours (Edwards *et al.*, 1977; Bruton *et al.*, 2008). This delayed force recovery at low frequencies has been termed prolonged low-frequency force depression (PLFFD).

Theoretically, PLFFD could be caused by an impaired SR Ca^{2+} release and/or reduced myofibrillar Ca^{2+} sensitivity. This is explained by the non-linear force- $[\text{Ca}^{2+}]_i$ relation, which is flat at high- while steep at low-stimulation frequencies. Hence, a reduction in tetanic $[\text{Ca}^{2+}]_i$ or a decrease in myofibrillar Ca^{2+} sensitivity will result in a more marked force reduction at low- than at high-frequencies (Westerblad *et al.*, 1993; Bruton *et al.*, 2008; Cheng *et al.*, 2015). The relative contribution of the two underlying causes of PLFFD varies with recovery time (Watanabe *et al.*, 2015; Watanabe & Wada, 2016) and experimental species (Bruton *et al.*, 2008).

Models to study skeletal muscle Ca^{2+} -handling, force and fatigue

Attempts to address the mechanisms behind skeletal muscle fatigue have employed experimental models that range from exercising humans to isolated single myofibrils (Lamb, 2002; Allen *et al.*, 2008). Given the central role of intracellular Ca^{2+} in the physiological events responsible for skeletal muscle force generation and fatigue development, experimental models that enable continuous recordings of $[\text{Ca}^{2+}]_i$ and force are best suited for such mechanistic studies. At the time this thesis work commenced, intact single fibres isolated from animal muscles were the only model to entirely fulfil these requirements (Place *et al.*, 2010). Due to the possibility of species differences, it was unknown to what extent findings in animal models would translate to the human setting. Alternative experimental models enabling studies of human skeletal muscle cells included *in vivo* studies, skinned skeletal muscle fibres, and cultures of myoblasts and myotubes. However, as will be alluded to next, inherent limitations of these models have limited their applicability in mechanistic studies of intracellular Ca^{2+} -handling and force production.

Human skeletal muscle *in vivo*

The intact human skeletal muscle *in vivo* with maintained blood supply, neural innervation of motor units and, often, mixture of fibre types, extracellular matrix and tendinous attachments intact is arguably the gold standard for experiments on human muscle functions. In the early 1950's, non-invasive experimental protocols

were developed to study human fatigue by analysing the reduction in maximal voluntary force following isometric contractions (Merton, 1954). Later, the possibility of EMG and percutaneous stimulation of the motor nerve or the muscle itself enabled the differentiation of peripheral and central components of human muscle fatigue (Bigland-Ritchie *et al.*, 1978). The muscle biopsy technique (Bergström, 1962), and non-invasive nuclear magnetic resonance spectroscopy (Sapega *et al.*, 1987), made it possible to study the correlation of skeletal muscle metabolic and ionic changes with fatigue development. Despite these advances, it remains difficult to distinguish between the different factors that may contribute to fatigue in human skeletal muscle *in vivo* experiments (Allen *et al.*, 2008). For these types of mechanistic studies, direct assessments of changes in $[Ca^{2+}]_i$ in parallel with measurements of force are generally considered necessary. Techniques that enable this type of studies in humans *in vivo* are yet to be developed.

Isolated intact single skeletal muscle fibres

Intact single skeletal muscle fibres can be isolated from animal muscles by either enzymatic dissociation or manual dissections. While the former represents a relatively straightforward procedure to obtain several muscle fibres per muscle bundle, the latter is a tedious dissection using forceps, micro-scissors and microscopes to obtain a single intact muscle fibre (Cheng & Westerblad, 2017). The great advantage of the latter approach is that, while the tendons are lost during enzymatic dissociation, they remain intact following manual dissection. This allows mounting of the muscle fibre in a force transducer making force recordings possible. Fluorescent Ca^{2+} indicator dyes can be introduced into viable muscle fibres, which enable continuous measurements of force and $[Ca^{2+}]_i$ in living muscle fibres (Place *et al.*, 2010; Cheng & Westerblad, 2017).

Skinned skeletal muscle fibres

Skinned skeletal muscle fibres are muscle fibres in which the sarcolemma has been disrupted or removed. This can be achieved either by chemical permeabilisation of the sarcolemma or by mechanically removing the sarcolemma (Lamb, 2002). The major advantage of the skinned muscle fibre model is that it does not require that intact muscle fibres be collected, which can be difficult to nearly impossible depending on the size of the muscle (Cheng & Westerblad, 2017). Notably, skinned muscle fibres can be obtained from, e.g., percutaneous muscle biopsies, enabling studies in human skeletal muscle (Place *et al.*, 2010).

Limitations of this model include a loss of soluble cytosolic proteins and the naturally occurring build-up of metabolic by-products that may be of importance in fatigue development. Skinned fibre experiments are generally conducted at temperatures lower than those prevailing *in vivo* and, as alluded to earlier, this may be problematic when considering the mechanistic aspects of fatigue (Lamb, 2002; Allen *et al.*, 2008). In addition, although feasible in the less used model of

mechanically skinned fibres (Posterino *et al.*, 2000), the standard procedure of chemically skinned fibres does not allow for analysis of SR Ca^{2+} release and excitation-contraction coupling (Lamb, 2002; Allen *et al.*, 2008).

Myoblasts and myotubes

Satellite cells are skeletal muscle resident stem cells involved in the maintenance and adaptive processes of skeletal muscle tissues (Mauro, 1961; Relaix & Zammit, 2012). Satellite cells can be extracted from human muscle biopsies, activated into proliferating myoblasts and differentiated into multinuclear myotubes *ex vivo* (Blau & Webster, 1981). Upon differentiation of myoblasts into myotubes, several morphological, metabolic and biochemical features similar to those in adult skeletal muscle fibres have been reported (Berggren *et al.*, 2007; Aas *et al.*, 2013). This together with the advantage that myotubes retain donor-specific traits have rendered myotubes a widely used *ex vivo* model of human skeletal muscle (Peter *et al.*, 2009; Ullrich *et al.*, 2011; Nikolic *et al.*, 2012; Aas *et al.*, 2013; Scheler *et al.*, 2013). However, differences in metabolic (Sarabia *et al.*, 1992; Baker *et al.*, 2003), contractile and intracellular Ca^{2+} -handling properties (Tanaka *et al.*, 2000; Bandi *et al.*, 2008; Rokach *et al.*, 2013; Cheng *et al.*, 2014; Smith *et al.*, 2014; Madden *et al.*, 2015) have been reported between myotubes and muscle fibres. These limitations cast doubt as to whether findings in myotubes are transferrable to the adult muscle fibre.

Intrinsic skeletal muscle changes in disease

Disease-related changes affecting skeletal muscle force producing capacity and fatigue resistance may occur at any level from the brain to the contractile proteins. Although likely the result of a combination of factors, it is clear that changes intrinsic to the skeletal muscle contributes to the loss of skeletal muscle function in patients suffering from chronic diseases including heart failure and rheumatoid arthritis (Middlekauff, 2010; Yamada *et al.*, 2017; Steinz *et al.*, 2019), but also in patients afflicted by acute critical illness myopathy (Friedrich *et al.*, 2015).

In heart failure patients, this is demonstrated by the finding of a remaining muscle force deficit when force is normalized to muscle cross sectional area, leg muscle mass, or body weight (Lipkin *et al.*, 1988; Harrington *et al.*, 1997; Toth *et al.*, 2006; Toth *et al.*, 2010; Toth *et al.*, 2012). A similar qualitative decline in muscle function is found in rheumatoid arthritis and critical illness myopathy patients (Helliwell & Jackson, 1994; Friedrich *et al.*, 2015), and rodent disease models (Yamada *et al.*, 2009; Yamada *et al.*, 2015a; Yamada *et al.*, 2015b; Llano-Diez *et al.*, 2016; Steinz *et al.*, 2019).

Intracellular Ca²⁺-handling and disease-related muscle dysfunction

Altered intracellular Ca²⁺-handling has been implicated in disease-related quantitative and qualitative changes in skeletal muscle. Changes in intracellular Ca²⁺-handling may involve principally two steps in the excitation-contraction coupling process that would directly affect muscle contractile function and resistance to fatigue: (i) the release/reuptake of SR Ca²⁺ and (ii) the myofibrillar Ca²⁺ sensitivity. Additionally, changes in basal [Ca²⁺]_i or [Ca²⁺]_i transient amplitude and/or duration regulate, e.g., muscle protein metabolism and the amount of contractile proteins available for force production (Bhattacharyya *et al.*, 1993; Menconi *et al.*, 2004; Smith & Dodd, 2007; Dridi *et al.*, 2020; Kushnir *et al.*, 2020).

A reduction in SR Ca²⁺ release and decreased myofibrillar Ca²⁺ sensitivity has been found to contribute to the muscle weakness observed in animal models of heart failure (Perreault *et al.*, 1993; Reiken *et al.*, 2003; Ward *et al.*, 2003), rheumatoid arthritis (Yamada *et al.*, 2009; Yamada *et al.*, 2015b), critical illness myopathy (Callahan *et al.*, 2001; Llano-Diez *et al.*, 2016), and normal ageing (Andersson *et al.*, 2011). Due to technical difficulties, mechanistic studies of this sort in humans are scarce. However, perturbed action potential-evoked Ca²⁺ transient amplitude and release in isolated skeletal muscle fibre segments from heart failure patients have been reported (DiFranco *et al.*, 2014). Additionally, the quantity and extent of posttranslational modifications of intracellular Ca²⁺-handling proteins differ between human heart failure patients and controls, indicating plausible differences in functional Ca²⁺-handling properties (Middlekauff *et al.*, 2012; Rullman *et al.*, 2013) in this patient group.

Causes of disease-related changes in intracellular Ca²⁺-handling

Studies in animal models have demonstrated that a loss of skeletal muscle mass and function, related to altered intracellular Ca²⁺-handling, is manifested in several diseases of otherwise different underlying aetiology (Reiken *et al.*, 2003; Yamada *et al.*, 2015b; Llano-Diez *et al.*, 2016). It is possible that factors not related to the disease process in itself, e.g., muscle disuse, or processes unique for each disease may explain these common manifestations. An alternative hypothesis is that the inherent biological response to several diseases shares common features, and that these contribute to the impaired muscle function. In support for this, several of these diseases exhibit common features, such as an increased neuroendocrine activation, elevated circulating levels of pro-

inflammatory cytokines, and altered sphingolipid metabolism (Andersson & Marks, 2010; Nikolova-Karakashian & Reid, 2011; Reid & Moylan, 2011). In animal models, these have all been demonstrated to induce changes in skeletal muscle intracellular Ca^{2+} -handling and to impair muscle function.

Neuroendocrine activation

Posttranslational modifications of RyR1, including phosphorylation, oxidation and nitrosylation, have been demonstrated in human heart failure patients (Rullman *et al.*, 2013). In animal models, these modifications have been shown to result in dissociation of the stabilising protein FKBP12 from RyR1, increasing the leakage of SR Ca^{2+} into the cytosol. This in turn causes an elevation in baseline $[\text{Ca}^{2+}]_i$ and a depletion of SR Ca^{2+} , ultimately causing a reduction in tetanic $[\text{Ca}^{2+}]_i$ amplitude and subsequent force decline (Ward *et al.*, 2003). Excessive neuroendocrine activation in heart failure has been suggested to cause skeletal muscle weakness and fatigue intolerance due to hyperphosphorylation of RyR1 via β -adrenergic activation (Reiken *et al.*, 2003).

Cytokines

A number of circulating proinflammatory cytokines are elevated in heart failure, rheumatoid arthritis and critical illness myopathy. Of these, TNF α is the cytokine most widely recognized to correlate with disease severity and poor prognosis (Rauchhaus *et al.*, 2000; Conraads *et al.*, 2003; Friedrich *et al.*, 2015). Albeit other circulating cytokines are elevated in these conditions, the correlation of these cytokines to muscle dysfunction and prognosis is weaker and more ambiguous (Testa *et al.*, 1996; Friedrich *et al.*, 2015). Increased levels of circulating TNF α , or soluble forms of TNF receptor I or II, which are believed to reflect an increased activity of TNF α , are elevated in heart failure (Levine *et al.*, 1990; Torre-Amione *et al.*, 1996; Rauchhaus *et al.*, 2000), and especially so in the most symptomatic heart failure patients (i.e. NYHA class III-IV) (Ferrari *et al.*, 1995; Testa *et al.*, 1996; Torre-Amione *et al.*, 1996; Rauchhaus *et al.*, 2000; Deswal *et al.*, 2001), and in cachectic patients (Levine *et al.*, 1990).

Exogenously administered TNF α depresses cardiomyocyte force by reducing tetanic $[\text{Ca}^{2+}]_i$ (Yokoyama *et al.*, 1993). In skeletal muscle, TNF α has been demonstrated to reduce membrane excitability suggesting that it may inactivate DHPR and thus reduce tetanic $[\text{Ca}^{2+}]_i$ and force (Tracey *et al.*, 1986). Later studies in isolated rodent skeletal muscle fibres have shown that TNF α acutely reduces tetanic force without affecting tetanic $[\text{Ca}^{2+}]_i$ (Reid *et al.*, 2002), by a RONS-dependent mechanism (Reid *et al.*, 2002; Stasko *et al.*, 2013). Notably, changes in intracellular Ca^{2+} -handling via RONS-induced modifications of the RyR1 have been suggested in human heart failure patients (Rullman *et al.*, 2013).

Sphingolipid metabolism and sphingomyelinase activity

The sphingomyelinase family is a group of biochemically and genetically different enzymes, all of which hydrolyse sphingomyelin to ceramide and phosphorylcholine (Nikolova-Karakashian & Reid, 2011). Circulating sphingomyelinase activity is elevated in human heart failure (Doehner *et al.*, 2007), rheumatoid arthritis (Hanaoka *et al.*, 2018) and critical illness myopathy patients (Claus *et al.*, 2005). In heart failure patients, circulating sphingomyelinase activity is positively correlated to both circulating TNF α and TNF-RI while negatively correlated to muscle strength measures and lean tissue mass (Doehner *et al.*, 2007). Moreover, circulating sphingomyelinase activity is increased in a stepwise fashion according to NYHA class while related to impaired survival, independent of age, NYHA class or mean blood pressure (Doehner *et al.*, 2007). Similarly, an elevation of sphingomyelinase activity predicts a worse outcome in patients treated at the intensive care unit, while inhibition of sphingomyelinase activity has been found to increase survival in a mouse model of endotoxic shock (Claus *et al.*, 2005).

In rodent skeletal muscle, exogenous sphingomyelinase increases ceramides and oxidant activity while perturbing force production (Ferreira *et al.*, 2010; Ferreira *et al.*, 2012; Loehr *et al.*, 2014; Bost *et al.*, 2015), and fatigue resistance (Ferreira *et al.*, 2010; Ferreira *et al.*, 2012; Loehr *et al.*, 2014). Sphingomyelinase-induced attenuation of force has been suggested to result from changes in maximum Ca²⁺-activated force and a reduction in myofibrillar Ca²⁺ sensitivity (Ferreira *et al.*, 2012). The current hypothesis is that increased levels of pro-inflammatory cytokines and RONS contribute to perturbed muscle function in diseases characterized by elevated sphingomyelinase activity via sphingomyelinase-dependent effects on skeletal muscle force producing capacity and fatigue resistance (Marathe *et al.*, 1998; Wong *et al.*, 2000; Jenkins *et al.*, 2010; Nikolova-Karakashian & Reid, 2011; Moylan *et al.*, 2014).

Disease-related loss of muscle mass

Disease-related loss of muscle mass results in a dismal life quality and has been reported as an independent risk factor for mortality (Anker *et al.*, 1997; Hulsmann *et al.*, 2004; Sokka *et al.*, 2008; Chung *et al.*, 2014; Powers *et al.*, 2016). The net protein balance, i.e. the ratio of protein synthesis and degradation, dictates muscle fibre size (Phillips *et al.*, 2012). When muscle protein degradation exceeds synthesis, muscle protein depletion ensues and the muscle fibre atrophies. At the whole tissue level, muscle atrophy may also be caused by the loss of individual muscle fibres (Wilkinson *et al.*, 2018). In several diseases, both a negative protein balance and a loss of muscle fibres have been shown to contribute to the loss of muscle mass (Powers *et al.*, 2016). Disease-related modifications of Ca²⁺-

regulated process have been postulated to trigger these changes (Bhattacharyya *et al.*, 1993; Menconi *et al.*, 2004; Smith & Dodd, 2007; Dridi *et al.*, 2020; Kushnir *et al.*, 2020).

A third potential cause of disease-related muscle atrophy is a reduced hypertrophic response and regenerative capacity of the skeletal muscle due to perturbed muscle stem cell function and abundance (McKenna & Fry, 2017; Wilkinson *et al.*, 2018). Muscle stem cell functions are regulated by local and systemic factors that modulate their proliferation and differentiation. One such factor that has increasingly been linked to skeletal muscle health and intimately linked to the regulation of muscle stem cell functions is vitamin D (Girgis *et al.*, 2013; Girgis *et al.*, 2014a).

The role of vitamin D in skeletal muscle physiology and pathology

Deficiency of Vitamin D is a widely spread and highly prevalent condition, and has been associated with skeletal muscle weakness and atrophy (Girgis *et al.*, 2013). The major circulating form of vitamin D consists of 25(OH)D, which is believed to be biologically inactive. 25(OH)D is hydroxylated by the enzyme 1 α -hydroxylase, present in the proximal tubules of the kidney, to form the biologically active 1 α ,25(OH)₂D. The half-life of 1 α ,25(OH)₂D is short, and the synthesis strictly controlled so that serum levels are usually well maintained despite very low levels of 25(OH)D (Need *et al.*, 2008). Assessment of vitamin D status is therefore made by measuring the serum concentration of 25(OH)D, upon which the diagnosis of vitamin D deficiency is made (Hollis & Wagner, 2005).

The vitamin D receptor has been reported to be present in virtually all cells of the body, including some reports claiming its presence in human skeletal muscle fibres (Pojednic *et al.*, 2015). Proposed effects of vitamin D in skeletal muscle include an increase in force generation via non-genomic actions on intracellular Ca²⁺-handling, and an anabolic response via transcriptional regulation (Girgis *et al.*, 2013; Bass *et al.*, 2020). In rodent muscles, overexpression of the vitamin D receptor induces a hypertrophic response by stimulating protein synthesis, and possibly muscle stem cell number and activity (Bass *et al.*, 2020). Notably, vitamin D has been suggested to reduce the expression of myostatin, a well-established negative regulator of muscle mass (McPherron *et al.*, 1997), in rodent skeletal muscle cells (Garcia *et al.*, 2011; Girgis *et al.*, 2014a). However, a direct effect of vitamin D on skeletal muscle cells has been a subject of controversy, with even the presence of the vitamin D receptor in the skeletal muscle tissue in question (Wang & DeLuca, 2011; Bouillon *et al.*, 2014; Pike, 2014).

2 Objectives and outline of the thesis

The overarching aim of the current thesis was to study the intracellular mechanisms of human muscle fatigue and disease-related loss of muscle function and mass. A special focus was put on the tentative role of altered intracellular Ca^{2+} -handling in these processes. An integrative approach spanning from single intact muscle fibres, isolated muscle stem cells, to whole muscle biopsies from healthy and diseased individuals was employed.

The specific aims of the studies were as follows:

- | | |
|------------------|--|
| <i>Study I</i> | To establish an experimental model suitable for mechanistic studies of intracellular Ca^{2+} -handling, force generation and fatigue in human skeletal muscle. |
| <i>Study II</i> | To test the hypothesis that vitamin D plays a role in maintaining human skeletal muscle function and mass by regulating muscle stem cell functions. |
| <i>Study III</i> | To investigate the cellular mechanisms of human muscle fatigue and the effects of acidosis on muscle performance. |
| <i>Study IV</i> | To test the hypothesis that elevated sphingomyelinase activity is implicated in disease-related loss of muscle function and mass by affecting intracellular Ca^{2+} -handling and muscle atrophy processes. |

3 Summary of study results

The following section provides a summary of the key results of studies I-IV. For detailed results the reader is referred to the individual studies presented in sections four to seven.

Study I: *Ex vivo* models of human skeletal muscle

In study I, we took advantage of the relatively short muscle fibres of the accessory respiratory muscles located between the ribs of humans, the intercostal muscles, to manually dissect intact single fibres. During thoracotomies performed for medical reasons, the intercostal muscles are exposed and dissected to access the thoracic cavity. Thus, they are accessible for muscle biopsies without unwarranted inconvenience or additional risk for the patient. By this approach we obtained intact single human muscle fibres that enabled electrically evoked contractions, while continuously measuring $[Ca^{2+}]_i$ and force. Human myoblasts, isolated from vastus lateralis muscle biopsies, were differentiated *ex vivo* to form myotubes, and their contractile function and Ca^{2+} -handling characteristics were compared to those of adult intercostal muscle fibres.

Upon differentiation, human myoblasts fused to form multinucleated myotubes with an increased expression of Ca^{2+} -handling proteins and functionality of the initial events of the excitation-contraction coupling cascade. Increased expression of contractile proteins and initiation of structural organisation were also evident upon differentiation. Yet, we found major remaining differences when compared to the adult muscle fibre: none of the myotubes contracted, and the Ca^{2+} transient decay upon electrical stimulation was markedly prolonged in myotubes (**Figure 3.1A-B**). Subsequent analyses suggested that these remaining differences were due to dissimilarities in regards to the protein abundance and co-localisation of Ca^{2+} -handling proteins, and ultrastructure (**Figure 3.1C-D**).

The duration of $[Ca^{2+}]_i$ transients have been demonstrated to affect the cellular signalling initiated by increased $[Ca^{2+}]_i$ (Berridge, 1990, 1992; Dolmetsch *et al.*, 1997; Berridge *et al.*, 2003; Clapham, 2007; Tavi & Westerblad, 2011). Thus, when studying Ca^{2+} -dependent processes such as gene expression and metabolism, myotubes are likely to provide different answers to those given by intact adult muscle fibres.




Figure 3.1. Human muscle fibres and myotubes showed marked differences in Ca^{2+} -handling characteristics and contractile ability. Changes in Ca^{2+} indicator dye intensity (A) and force (B) following electrical stimulation of a human muscle fibre, myotube and myoblast. Fluorescent intensity for RyR and DHPR (C), and TEM images of the ultrastructure (D) in a human muscle fibre and myotube. Note the characteristic arrangement of myofibrils separated by triads (arrow) at the junction of the A and I bands in fibres while only rudimentary myofilament structures (*) were seen in myotubes.

Study II: Effects of vitamin D in human skeletal muscle

Contemplating previous findings and the results in study I, we considered human myoblasts, myotubes and muscle fibres to represent distinct stages of muscle development. In study II, we took advantage of this to investigate the presence of components of the vitamin D endocrine system, including the vitamin D receptor, and the vitamin D metabolising enzymes CYP24A1 and CYP27B1, in human muscle cells at different developmental stages. The effects of vitamin D on the regulation of processes involved in muscle regeneration and growth were investigated in human myoblasts.

The vitamin D receptor was readily detectable in human myoblasts and myotubes, while non-detectable in human adult skeletal muscle (**Figure 3.2A**). Attention was therefore focused on the effects of vitamin D on human myoblast proliferation and differentiation. Results revealed a profound inhibitory effect by the biologically active form of vitamin D ($1\alpha,25(\text{OH})_2\text{D}$) on myoblast proliferation and differentiation. No effects by the biologically inactive form $25(\text{OH})_2\text{D}$ were detected in human myoblasts (**Figure 3.2B**).




Figure 3.2. The vitamin D receptor was readily detected in human muscle stem cells and vitamin D markedly inhibited human myotube formation. A, protein quantity of the vitamin D receptor (VDR) in human primary myoblasts, myotubes and adult skeletal muscle. B, fusion index following 8 days of differentiation in the presence of 1 nM or 100 nM $1\alpha,25(\text{OH})_2\text{D}_3$, or 100 nM $25(\text{OH})\text{D}_3$. Data are presented as the mean \pm SEM. * $P < 0.05$, ** $P < 0.01$. (ND) = non-detectable.

Global gene expression analysis provided mechanistic insight by demonstrating an altered expression of cell cycle regulators consistent with a block of cell cycle progression at the G_1/S restriction point. Further, vitamin D reduced the expression of MyoD, myogenin and MEF2C, myogenic regulatory factors acknowledged for their role in muscle stem cell commitment and differentiation (Relaix & Zammit, 2012). Gene-ontology enrichment of down-regulated genes identified GO categories involved in ‘muscle organ development’. Network analysis using IPA identified pathways annotated as ‘differentiation of myoblasts’ and ‘differentiation of muscle cells’ among the most highly inhibited biological functions. Altered expression of genes intimately involved in muscle stem cell self-renewal lead us to suggest a role of the biologically active form of vitamin D in the maintenance of the muscle stem cell pool (**Figure 3.3**).




Figure 3.3. Illustration of the results from studies I and II. Vitamin D promotes satellite cell self-renewal while inhibiting myoblast differentiation. Activated myoblasts form myotubes both *in vivo* and *ex vivo*, but only *in vivo* do myotubes fully mature into muscle fibres with functional excitation-contraction (E-C) coupling and contractile ability.

Study III: Cellular mechanisms of human skeletal muscle fatigue

In study III, the cellular mechanisms of human muscle fatigue following near maximal isometric contractions were investigated by continuous measurements of force and $[Ca^{2+}]_i$ transients. Based on their ability to maintain force >40% of the unfatigued value during 500 tetani, two distinct groups of fibres were distinguished: fatigue sensitive and fatigue resistant.

In both groups of fibres, the fatigue-induced force loss was primarily caused by an impaired SR Ca^{2+} release, while the myofibrillar Ca^{2+} sensitivity was either unaltered (fatigue-sensitive fibres) or increased (fatigue-resistant fibres) (**Figure 3.4**). Integrating these results, fatigue development in human muscle fibres is different to that in rodent muscle fibres in that reduced myofibrillar Ca^{2+} sensitivity plays no role in the force loss in human fibres.




Figure 3.4. Illustration of the cellular mechanisms of fatigue in human muscle fibres. A and B, contractile-induced changes in tetanic force, $[Ca^{2+}]_i$ and force- $[Ca^{2+}]_i$ relationship in human fatigue-sensitive and fatigue-resistant muscle fibres, respectively. Note that in both human fatigue-sensitive and fatigue-resistant fibres the main cause of fatigue was a reduced SR Ca^{2+} release while myofibrillar Ca^{2+} sensitivity was either unaltered (fatigue-sensitive fibres) or increased (fatigue-resistant fibres) following fatigue.

In fatigue-resistant fibres, recovery was complete 15 minutes after cessation of fatiguing contractions. In contrast, in fatigue-sensitive fibres the force at low-frequency stimulation was markedly lower, while force at high-frequency stimulation was recovered, compared to the unfatigued value. This phenomenon, known as prolonged low-frequency force depression (PLFFD), was exclusively due to an impaired SR Ca^{2+} release. Myofibrillar Ca^{2+} sensitivity and maximal force producing capacity was unaltered (**Figure 3.5**). Cumulatively, these results demonstrate a fundamental importance of impaired SR Ca^{2+} release to the force depression following fatigue and in the early recovery period in human muscle fibres.




Figure 3.5. PLFFD in human muscle fibres was caused exclusively by an impaired SR Ca^{2+} release. A, representative force and $[\text{Ca}^{2+}]_i$ records in a fatigue-sensitive muscle fibre at low- and high-frequency stimulation before (unfatigued) and 15 min after (recovery) fatiguing stimulation. C, PLFFD was evident in human fatigue-sensitive muscle fibres. D, the force decline in fatigue-sensitive fibres during recovery followed the unfatigued force- $[\text{Ca}^{2+}]_i$ relationship with force fully restored to unfatigued levels in the presence of 5 mM caffeine. Data are means \pm SEM. * $P < 0.05$, ** $P < 0.01$.

Intracellular acidosis has historically been ascribed a major causative role in the fatigue-induced force loss following intense contractile activity (Fitts, 2016). However, as alluded to previously, this view has later been challenged by findings in animal isolated muscles studied at or near physiological temperatures (Pate *et al.*, 1995; Westerblad *et al.*, 1997; Bruton *et al.*, 1998). We were therefore interested in the effects of intracellular acidosis on the force- $[\text{Ca}^{2+}]_i$ relation and fatigue resistance of human fibres at 37° C. In unfatigued human muscle fibres, intracellular acidosis reduced the myofibrillar Ca^{2+} sensitivity, but this was completely offset by an increase in tetanic $[\text{Ca}^{2+}]_i$ so that force was maintained (**Figure 3.6A-B**). Results further suggested that the fatigue-resistance of human fibres was unaltered upon acidification (**Figure 3.6C-D**).




Figure 3.6. Acidosis did not reduce force production, nor decrease the fatigue tolerance of human intercostal muscle fibres. A, representative $[Ca^{2+}]_i$ and force records in a human muscle fibre during a 30 Hz and 70 Hz tetanus under control (5% CO_2) and acidic (30% CO_2) conditions. B, acidification resulted in increased tetanic $[Ca^{2+}]_i$ while force production was unaltered. C and D, number of tetanic contractions required to reach 40% of the initial force in fatigue-sensitive (C) and fatigue-resistant (D) human fibres. Data are means with individual data points as well as the 90% confidence interval of the group mean difference (C and D). Adjoining lines indicate data points from paired experiments. * $P < 0.05$, ** $P < 0.01$.

Study IV: The role of sphingomyelinase in disease-related loss of muscle function and mass

In study IV, the effects of sphingomyelinase on the force production, Ca^{2+} -handling, and transcriptional regulation of genes related to muscle atrophy processes were studied in intact muscle fibres. The activity of sphingomyelinase was measured in skeletal muscle from human heart failure and healthy age-matched subjects, and related to circulatory markers of inflammation and factors of known prognostic value in heart failure, previously reported by our lab for the current patient cohort (Rullman *et al.*, 2020).

The results show that sphingomyelinase activity impaired SR Ca^{2+} release and reduced myofibrillar Ca^{2+} sensitivity to depress force in intact human muscle fibres (**Figure 3.7**).




Figure 3.7. Illustration of the effects of sphingomyelinase on muscle fibre force and Ca^{2+} -handling. Sphingomyelinase-induced changes in tetanic force and $[\text{Ca}^{2+}]_i$ (A) and in the force- $[\text{Ca}^{2+}]_i$ relationship (B) in human intact muscle fibres. Sphingomyelinase activity reduced myofibrillar Ca^{2+} sensitivity (i) and decreased tetanic $[\text{Ca}^{2+}]_i$ (ii). SMase, sphingomyelinase.

In human muscle bundles, sphingomyelinase induced the transcription of genes of the ubiquitin-proteasome system, including essential subunits of the proteasome 20s core (PSMB5, 6, 7 and PSMA4), the 19s regulator (PSMC 1, 3, 4, 5 and PSMD1), polyubiquitin B and C (**Figure 3.8A**). Meanwhile, transcripts of ribosomal proteins (**Figure 3.8B**), factors implicated in muscle proliferation, and cellular locomotion were reduced by sphingomyelinase.




Figure 3.8. Sphingomyelinase activity induced transcriptional changes promoting muscle atrophy in human intact muscle fibres. Barcode-plots showing sphingomyelinase-induced changes in factors involved in proteasome degradation (A) and ribosomal proteins (B). In the lower panel each regulated gene of interest is depicted as a vertical line on the corresponding ratio (blue panel depicts down-regulated and red up-regulated genes) in treated vs. control-muscle. The upper panel depicts the enrichment of fold-changes between treated vs. control-muscle.

In human heart failure patients, the activity of intramuscular neutral sphingomyelinase (**Figure 3.9A**), but not acid sphingomyelinase, was increased as compared to age-matched healthy controls. In the cohort of patients included in the longitudinal axis of the study, both neutral (**Figure 3.9B**) and acid sphingomyelinase activity were found to increase with duration of disease. Intramuscular neutral sphingomyelinase activity positively correlated with circulatory markers of inflammatory activation and atrophy including TNF-RI and II, and markers previously demonstrated to carry prognostic value in this patient cohort (Rullman *et al.*, 2020), including NT-proBNP, IGFBP7 and GDF15 (**Figure 3.9C**).




Figure 3.9. Intramuscular neutral sphingomyelinase activity is elevated in human heart failure and correlates to circulatory markers of inflammation and muscle atrophy. A, skeletal muscle nSMase activity in human heart failure patients and healthy age-matched controls. C, skeletal muscle nSMase activity in human heart failure patients at the time of the baseline biopsy (time of inclusion in the study) and at the follow-up biopsy. Mean follow-up time $2.5 (\pm 0.2)$ years. Bars indicate mean value and circles represent individual data points. C, correlation of skeletal muscle nSMase activity and serum protein quantities of factors analysed with targeted proteomics in human heart failure patients ($n = 61$) where edges denote correlation coefficients >0.3 . nSMase, neutral sphingomyelinase.

8 Discussion and implications

This thesis investigated the cellular mechanisms underlying human muscle fatigue and disease-related muscle dysfunction, with a special focus on Ca^{2+} -regulated processes. Novel contributions include a mechanistic insight into the underlying causes of fatigue in human muscle fibres, the direct effects of vitamin D in muscle, and the role of sphingomyelinase in disease-induced muscle weakness and atrophy.

Ex vivo models of human skeletal muscle

At the time this thesis work commenced, manually dissected single fibres were arguable still the best model for mechanistic studies of muscle fibre functions. An obvious drawback of this model is the inferred problem of species differences, i.e., can we be sure that the findings in rodent muscles are representative of those in human muscles? A tentative approach to overcome this issue is the use of differentiated human myotubes. These can be obtained from minimally invasive muscle biopsies (Blau & Webster, 1981), and have been acclaimed traits that are similar to muscle fibres. In addition, they have been shown to retain donor-specific qualities (Berggren *et al.*, 2007; Aas *et al.*, 2013). The possibility to study disease-related changes in human muscle cells spurred our interest to investigate the utility of myotubes for studies of muscle fibre functions.

In study I, myotubes were found to develop some fibre-like qualities, but they did not develop functional excitation-contraction coupling. Hence, at least using the 2D culture protocol employed in study I, myotubes were found to be an inadequate model for studies of these muscle fibre functions. At the time of study I and in the years that followed, 3D cultures and continuous electrical stimulation were demonstrated to ameliorate the development of functional excitation-contraction coupling in myotubes (Madden *et al.*, 2015; Rao *et al.*, 2018; Khodabukus *et al.*, 2019). Despite this, major differences in force production and Ca^{2+} transient kinetics were still apparent between these bioengineered myotubes and muscle fibres. Manually dissected single fibres from human intercostal muscles were therefore chosen in the following studies of adult muscle fibre functions in this thesis. However, the technical difficulties related to the dissection and the dependence of surgical procedures makes this a time-consuming and low-throughput model. Continuous efforts to develop *ex vivo* models of human myotubes amenable for studies of muscle fibre contraction and Ca^{2+} -handling are therefore anticipated and warranted.

Effects of vitamin D in human skeletal muscle

In study II, we found components of the vitamin D endocrine system were abundant in human muscle stem cells, while low to non-detectable in adult skeletal muscle. The apparent lack of the VDR in adult muscle fibres contrasts some previous reports in human and rodent skeletal muscle (Simpson *et al.*, 1985; Costa *et al.*, 1986; Bischoff *et al.*, 2001; Pojednic *et al.*, 2015). The reason for this disparity is likely due to differences in the antibody used to detect the VDR. Notably, when using the stringently validated VDR D-6 antibody, Wang and DeLuca (2011) reported that the VDR was un-detectable in skeletal muscle fibres, while confirming unspecific binding of other commercially available antibodies (Wang & DeLuca, 2011). While this does not exclude a direct role of vitamin D in muscle fibres, these findings suggest direct effects of vitamin D in human skeletal muscle to predominantly involve muscle stem cells.

Coherent with such a role, results in study II demonstrated a robust direct effect of vitamin D in human muscle stem cells: the biologically active $1\alpha,25(\text{OH})_2\text{D}_3$, but not $25(\text{OH})\text{D}_3$, inhibited human myoblast proliferation and differentiation. This marks a contribution to the literature, as previous studies have been limited to animal muscle stem cells (Garcia *et al.*, 2011; Srikuea *et al.*, 2012; Girgis *et al.*, 2014a; Girgis *et al.*, 2014b).

In order to maintain the skeletal muscle stem cell pool, part of the myoblast progeny must return to quiescence by events inhibiting myoblast proliferation and differentiation (Kuang *et al.*, 2007). This process is controlled by the FOXO3 and Notch signalling pathways (Kuang *et al.*, 2007; Bjornson *et al.*, 2012; Gopinath *et al.*, 2014), both of which we found to be modulated by vitamin D. Corroborating such an action, vitamin D deficiency has been suggested to contribute to the advent of sarcopenia in rodent muscles via diminished muscle stem cell proliferative and skeletal muscle regenerative potential by down-regulation of the Notch signalling pathway (Domingues-Faria *et al.*, 2014). Notably, increased fibre size and strength has been reported in rat muscles electroporated with the vitamin D receptor (Bass *et al.*, 2020). These effects were attributed partly to an increase in muscle stem cell activity.

Collectively, our results suggest that vitamin D plays a role in muscle stem cell self-renewal and quiescence, possibly by modulating the FOXO3 and Notch signalling pathways. This could be of importance for the overall functionality of the skeletal muscle tissue, especially in the ageing individual. The levels of the biologically active form of vitamin D is strictly regulated and well maintained despite very low levels of the prohormone $25(\text{OH})\text{D}$ (Need *et al.*, 2008). Impaired skeletal muscle function due to vitamin D deficiency is therefore likely uncommon. However, this will be of importance in conditions associated with a decline in the synthesis of the biologically active form of vitamin D, such as in chronic kidney disease, or in severe vitamin D deficiency.

Cellular mechanisms of fatigue in human muscle fibres

Study III highlighted that impaired SR Ca^{2+} release has a prominent role in the force loss during fatigue and in the early recovery period in human muscle fibres. This was in line with previous papers identifying reduced SR Ca^{2+} release as an important underlying cause of muscle fatigue in rodent fibres (Westerblad & Allen, 1991, 1993a). Impaired SR Ca^{2+} release is believed to stem from Ca^{2+} - P_i precipitation in the SR (Fryer *et al.*, 1995; Westerblad & Allen, 1996), in combination with an inhibition of the RyR1 channel caused by the depletion of cytosolic [ATP] and accumulation of free $[\text{Mg}^{2+}]$ (Blazev & Lamb, 1999).

A distinct initial phase of increased tetanic $[\text{Ca}^{2+}]_i$ while tetanic force declines, previously described in rodent fibres, was observed in human fatigue-sensitive fibres. This early force reduction has been attributed primarily to accumulation of P_i that depresses maximal Ca^{2+} -activated force (Pate & Cooke, 1989; Millar & Homsher, 1990; Dahlstedt *et al.*, 2000). Considering the identical pattern seen in fatigue-sensitive human fibres, we suggest this to be the most likely cause also in human fibres.

Unlike in rodent fibres, a reduction in myofibrillar Ca^{2+} sensitivity did not contribute to fatigue in human muscle fibres. Intriguingly, an increase in myofibrillar Ca^{2+} sensitivity was seen in fatigue-resistant fibres. This implies that changes in myofibrillar Ca^{2+} sensitivity can contribute to the fatigue-resistance of slow-twitch human fibres. The underlying mechanisms of fatigue between human and rodent muscle fibres are illustrated in **Figure 8.1**.




Figure 8.1. Illustration of the cellular mechanisms of fatigue in rodent and human muscle fibres. Contribution of changes in maximal Ca^{2+} -activated force, reduced SR Ca^{2+} release, and decreased myofibrillar Ca^{2+} sensitivity to fatigue in rodent (A) and human (B) fast-twitch fibres. Note that in rodent fibres (A) both a reduced SR Ca^{2+} release and decreased myofibrillar Ca^{2+} sensitivity contributes to the accelerated final decline in force, while in human fibres (B) this is caused exclusively by an impaired SR Ca^{2+} release. Force- $[\text{Ca}^{2+}]_i$ relation in unfatigued (solid line) and fatigued (dashed line) fibres.

Fatigue-induced changes in myofibrillar Ca^{2+} sensitivity have been investigated in human skinned fibres subjected to prior fatiguing stimulation *in vivo*. Results from these studies suggest that contractile-induced changes are multifaceted and can induce both an increased and decreased myofibrillar Ca^{2+} sensitivity of slow- and fast-twitch fibres (Hvid *et al.*, 2013; Gejl *et al.*, 2016; Lambole *et al.*, 2020). Contractile-induced RONS is one tentative factor involved in the modulation of myofibrillar Ca^{2+} sensitivity. Notably, RONS-mediated modifications of the fast troponin I isoform have been linked to both an increase and a decrease in myofibrillar Ca^{2+} sensitivity (Murphy *et al.*, 2008; Mollica *et al.*, 2012; Dutka *et al.*, 2017). This was explained by a competitive action on the Cys134 residue of troponin I by different RONS-species, which could explain the reported diversity of changes in myofibrillar Ca^{2+} sensitivity. Additionally, phosphorylation of the regulatory myosin light chain of fast-twitch fibres has been acknowledged to increase myofibrillar Ca^{2+} sensitivity (Sweeney *et al.*, 1993). An alternative hypothesis is that ADP and AMP accumulates in human fatigue-resistant fibres, due to the lower AMP deaminase and adenylate kinase activity in slow-twitch than in fast-twitch fibres (Borges & Essén-Gustavsson, 1989; Norman *et al.*, 1994). Notably, ADP has been shown to increase the myofibrillar Ca^{2+} sensitivity of slow-twitch fibres (Macdonald & Stephenson, 2006).

Acidosis and human muscle fatigue

Coherent with findings in rodents, intracellular acidosis caused a decrease in myofibrillar Ca^{2+} sensitivity while tetanic $[\text{Ca}^{2+}]_i$ increased in unfatigued human fibres. These changes inferred under acidic conditions in unfatigued fibres are strikingly different to the fatigue-induced changes under control conditions. This indicates that intracellular acidosis does not occur during fatigue under the present conditions or, alternatively, that other contractile-induced changes completely offset the effects of acidosis. Owing to a lack of tissue, fatigue-induced changes in pH_i were not measured in any of the human fibres. Therefore, we cannot exclude the possibility that fatigue under the current experimental conditions results in a less marked drop in pH_i than what is seen *in vivo*. Notably, in rodent fibres fatigued by similar experimental protocols as in the present study, development of intracellular acidosis of a magnitude similar to that in human muscles fatigued *in vivo* has been reported (Sahlin *et al.*, 1975; Sahlin *et al.*, 1976; Bruton *et al.*, 1998). Major differences in fatigue-induced changes in pH_i between the current experimental setting and the *in vivo* situation thus appear unlikely. To fully address these questions, future studies should include measurements of fatigue-induced changes in pH_i and $[\text{Ca}^{2+}]_i$ under control and acidic conditions, respectively.

In order to maintain tetanic force following a decrease in myofibrillar Ca^{2+} sensitivity, tetanic $[\text{Ca}^{2+}]_i$ must be adjusted accordingly. This will infer an increased energetic cost of force production due to the demand for energy-dependent removal of Ca^{2+} during relaxation. In analogy, fatigue is mitigated when myofibrillar Ca^{2+} sensitivity is increased (Cheng *et al.*, 2019). It thus appears likely that acidosis would exacerbate fatigue by increasing the metabolic cost of force production. This may become evident when combined with other mechanisms affecting force production, as previously demonstrated under circumstances of severe acidosis combined with markedly elevated P_i concentrations (Cooke *et al.*, 1988; Nelson *et al.*, 2014; Nelson & Fitts, 2014; Sundberg *et al.*, 2018). Repeated sprints may be such an example where severe acidosis is combined with profound changes in metabolites to further aggravate fatigue (**Figure 8.2**).




Figure 8.2. Tentative effects of acidosis on human muscle fatigue tolerance. A, the myofibrillar Ca^{2+} sensitivity is reduced under acidic conditions, but this effect is offset by an increase in tetanic $[\text{Ca}^{2+}]_i$ in unfatigued human muscle fibres. B and C, illustration of the hypothetical response to repeated fatigue bouts in human muscle fibres under non-acidic (B) and acidic conditions (C).

The role of sphingomyelinase in disease-related loss of muscle function and mass

Study IV implicate elevated sphingomyelinase activity as an underlying cause of disease-related loss of muscle function and mass. The following findings support this notion: (i) sphingomyelinase decreased myofibrillar Ca^{2+} sensitivity and impaired SR Ca^{2+} release to depress force in intact muscle fibres, (ii) transcriptional changes promoting muscle atrophy processes were induced upon sphingomyelinase exposure in human fibres, (iii) the activity of intramuscular sphingomyelinase was elevated in heart failure compared to control individuals and (iv) positively correlated with circulating markers of inflammation, atrophy and factors carrying prognostic value in heart failure.

The effects of sphingomyelinase on force generation and myofibrillar Ca^{2+} sensitivity are in line with previous reports in rodent fibres (Ferreira *et al.*, 2012; Empinado *et al.*, 2014; Loehr *et al.*, 2014; Bost *et al.*, 2015). We report the first study of the effects of sphingomyelinase on SR Ca^{2+} release. Previous studies have been limited to chemically skinned fibres and muscle bundles, which do not allow such investigations (Place *et al.*, 2010; Cheng & Westerblad, 2017). By demonstrating a depressive effect of sphingomyelinase on SR Ca^{2+} release, our data thus add novel insight into the force depression caused by sphingomyelinase.

In daily life, the ability to endure a physical activity at a set intensity or load over time (i.e. fatigue resistance) is dictated by the muscle strength and endurance. With all other variables set, if muscle strength declines this will demand that each contraction requires a higher α -motoneuron firing frequency in order to elevate $[\text{Ca}^{2+}]$, and to produce the required force. This will increase the energetic cost of each contraction and inevitably shorten the time to task failure, i.e. exhaustion. The muscle weakness caused by sphingomyelinase will thus be experienced as a decreased fatigue resistance in afflicted patients (**Figure 8.3**). In severe cases, the demands of daily life activities (e.g. rising up from a chair, walking up stairs) will demand near maximal Ca^{2+} -activated contractions and will be perceived as profound and debilitating muscle weakness and fatigue intolerance.

In human muscle fibres, transcriptional changes induced by sphingomyelinase included an up-regulation of genes involved in proteasome degradation and a down-regulation of genes encoding ribosomal proteins. The ubiquitin-proteasome is one of the two major proteolytic systems reported to be involved in muscle atrophy (Bonaldi & Sandri, 2013). Gene expression analyses of diverse clinical conditions associated with muscle atrophy have revealed common transcriptional changes of genes regulating the ubiquitin-proteasome system (Bodine *et al.*, 2001; Gomes *et al.*, 2001). These include an up-regulation of transcripts encoding ubiquitin, ubiquitin-conjugating enzymes, ubiquitin-protein ligases and several proteasome subunits (Lecker *et al.*, 2004). Notably, several of these common

transcripts, including essential subunits of the 20s core and 19s regulatory proteasome components, as well as polyubiquitin C, were induced in human muscle fibres exposed to sphingomyelinase. Among the ribosomal proteins transcriptionally repressed following sphingomyelinase exposure were RPL6, a component of the 60s ribosomal subunit also known to be an intracellular partner of FGF2 (Shen *et al.*, 1998). RPL6-FGF2 interaction is thought to promote cell survival (Sluzalska *et al.*, 2020) and thus suppressed RPL6 transcription may promote atrophy via cellular apoptosis. In light of the reported association of sphingomyelinase activity and muscle atrophy, e.g., in heart failure patients (Doehner *et al.*, 2007), and our finding of elevated sphingomyelinase activity in skeletal muscle of heart failure patients, a decisive role of sphingomyelinase in disease-related muscle atrophy is suggested.

In skeletal muscle, TNF α acts through TNF-R1 to activate neutral sphingomyelinase-3 (Moylan *et al.*, 2014). Neutral sphingomyelinase-3 does not have a well-defined role in other organs and does not appear to be critical for general sphingolipid metabolism. Hence, it provides a tentative target for selective inhibition that could mitigate disease-related muscle dysfunction and muscle wasting. Given the emerging role of sphingomyelinase activation in various clinical conditions, and the recent introduction of inhibitors of other sphingomyelinase isoforms (Sala *et al.*, 2020), this will be an interesting area of research to follow in the future.




Figure 8.3. Illustration of the effects of sphingomyelinase on human muscle force production, tetanic [Ca²⁺]_i, and plausible effects on fatigue tolerance. A, sphingomyelinase depresses force by reducing tetanic [Ca²⁺]_i and decreasing myofibrillar Ca²⁺ sensitivity. B, plausible effects of intracellular sphingomyelinase on the fatigue tolerance of human muscle fibres. A and B, sphingomyelinase reduces myofibrillar Ca²⁺ sensitivity so that larger increases in [Ca²⁺]_i are needed to attain maximal Ca²⁺-activated force (i), while equimolar [Ca²⁺]_i produces less force than in controls (ii). The combined effects of reduced myofibrillar Ca²⁺ sensitivity and impaired SR Ca²⁺ release severely depresses force in fibres exposed to sphingomyelinase (iii).

References

- Aas V, Bakke SS, Feng YZ, Kase ET, Jensen J, Bajpeyi S, Thoresen GH & Rustan AC. (2013). Are cultured human myotubes far from home? *Cell Tissue Res* **354**, 671-682.
- Allen DG, Lamb GD & Westerblad H. (2008). Skeletal muscle fatigue: cellular mechanisms. *Physiol Rev* **88**, 287-332.
- Allen DG, Whitehead NP & Froehner SC. (2016). Absence of Dystrophin Disrupts Skeletal Muscle Signaling: Roles of Ca^{2+} , Reactive Oxygen Species, and Nitric Oxide in the Development of Muscular Dystrophy. *Physiol Rev* **96**, 253-305.
- Andersen CL, Jensen JL & Ørntoft TF. (2004). Normalization of real-time quantitative reverse transcription-PCR data: a model-based variance estimation approach to identify genes suited for normalization, applied to bladder and colon cancer data sets. *Cancer Res* **64**, 5245-5250.
- Andersson DC, Betzenhauser MJ, Reiken S, Meli AC, Umanskaya A, Xie W, Shiomi T, Zalk R, Lacampagne A & Marks AR. (2011). Ryanodine receptor oxidation causes intracellular calcium leak and muscle weakness in aging. *Cell Metab* **14**, 196-207.
- Andersson DC & Marks AR. (2010). Fixing ryanodine receptor Ca^{2+} leak - a novel therapeutic strategy for contractile failure in heart and skeletal muscle. *Drug Discov Today Dis Mech* **7**, e151-e157.
- Andrade FH, Reid MB, Allen DG & Westerblad H. (1998). Effect of hydrogen peroxide and dithiothreitol on contractile function of single skeletal muscle fibres from the mouse. *J Physiol* **509** (Pt 2), 565-575.
- Anker SD, Ponikowski P, Varney S, Chua TP, Clark AL, Webb-Peploe KM, Harrington D, Kox WJ, Poole-Wilson PA & Coats AJ. (1997). Wasting as independent risk factor for mortality in chronic heart failure. *Lancet* **349**, 1050-1053.
- Baker EL, Dennis RG & Larkin LM. (2003). Glucose transporter content and glucose uptake in skeletal muscle constructs engineered in vitro. *In Vitro Cell Dev Biol Anim* **39**, 434-439.
- Bandi E, Jevsek M, Mars T, Jurdana M, Formaggio E, Sciancalepore M, Fumagalli G, Grubic Z, Ruzzier F & Lorenzon P. (2008). Neural agrin controls maturation of the excitation-contraction coupling mechanism in human myotubes developing in vitro. *Am J Physiol Cell Physiol* **294**, C66-73.
- Bangsbo J, Madsen K, Kiens B & Richter EA. (1996). Effect of muscle acidity on muscle metabolism and fatigue during intense exercise in man. *J Physiol* **495** (Pt 2), 587-596.
- Bass JJ, Nakhuda A, Deane CS, Brook MS, Wilkinson DJ, Phillips BE, Philp A, Tarum J, Kadi F, Andersen D, Garcia AM, Smith K, Gallagher IJ, Szewczyk NJ, Cleasby ME & Atherton PJ. (2020). Over-expression of the vitamin D receptor (VDR) induces skeletal muscle hypertrophy. *Mol Metab*, 101059.

- Beaudart C, Buckinx F, Rabenda V, Gillain S, Cavalier E, Slomian J, Petermans J, Reginster JY & Bruyere O. (2014). The effects of vitamin D on skeletal muscle strength, muscle mass, and muscle power: a systematic review and meta-analysis of randomized controlled trials. *J Clin Endocrinol Metab* **99**, 4336-4345.
- Bentzinger CF, Wang YX & Rudnicki MA. (2012). Building muscle: molecular regulation of myogenesis. *Cold Spring Harb Perspect Biol* **4**.
- Berchtold MW, Brinkmeier H & Muntener M. (2000). Calcium ion in skeletal muscle: its crucial role for muscle function, plasticity, and disease. *Physiol Rev* **80**, 1215-1265.
- Berggren JR, Tanner CJ & Houmard JA. (2007). Primary cell cultures in the study of human muscle metabolism. *Exerc Sport Sci Rev* **35**, 56-61.
- Bergström J. (1962). Muscle electrolytes in man. *Scand J Clin Lab Invest* **14**, 1-110.
- Berridge MJ. (1990). Temporal aspects of calcium signalling. *Adv Second Messenger Phosphoprotein Res* **24**, 108-114.
- Berridge MJ. (1992). Spatiotemporal aspects of calcium signalling. *Jpn J Pharmacol* **58 Suppl 2**, 142P-149P.
- Berridge MJ, Bootman MD & Roderick HL. (2003). Calcium signalling: dynamics, homeostasis and remodelling. *Nat Rev Mol Cell Biol* **4**, 517-529.
- Bhattacharyya J, Thompson KD & Sayeed MM. (1993). Skeletal muscle Ca^{2+} flux and catabolic response during sepsis. *Am J Physiol* **265**, R487-493.
- Bigland-Ritchie B, Jones DA, Hosking GP & Edwards RH. (1978). Central and peripheral fatigue in sustained maximum voluntary contractions of human quadriceps muscle. *Clin Sci Mol Med* **54**, 609-614.
- Bischoff HA, Borchers M, Gudat F, Duermueller U, Theiler R, Stahelin HB & Dick W. (2001). In situ detection of 1,25-dihydroxyvitamin D_3 receptor in human skeletal muscle tissue. *Histochem J* **33**, 19-24.
- Bjornson CR, Cheung TH, Liu L, Tripathi PV, Steeper KM & Rando TA. (2012). Notch signaling is necessary to maintain quiescence in adult muscle stem cells. *Stem cells* **30**, 232-242.
- Blau HM & Webster C. (1981). Isolation and characterization of human muscle cells. *Proc Natl Acad Sci U S A* **78**, 5623-5627.
- Bodine SC, Latres E, Baumhueter S, Lai VK, Nunez L, Clarke BA, Poueymirou WT, Panaro FJ, Na E, Dharmarajan K, Pan ZQ, Valenzuela DM, DeChiara TM, Stitt TN, Yancopoulos GD & Glass DJ. (2001). Identification of ubiquitin ligases required for skeletal muscle atrophy. *Science* **294**, 1704-1708.
- Bonaldo P & Sandri M. (2013). Cellular and molecular mechanisms of muscle atrophy. *Dis Model Mech* **6**, 25-39.
- Borges O & Essén-Gustavsson B. (1989). Enzyme activities in type I and II muscle fibres of human muscle in relation to age and torque development. *Acta Physiol Scand* **136**, 29-31.

- Bost ER, Frye GS, Ahn B & Ferreira LF. (2015). Diaphragm dysfunction caused by sphingomyelinase requires the p47^(phox) subunit of NADPH oxidase. *Respir Physiol Neurobiol* **205**, 47-52.
- Bouillon R, Gielen E & Vanderschueren D. (2014). Vitamin D receptor and vitamin D action in muscle. *Endocrinology* **155**, 3210-3213.
- Bruton J, Tavi P, Aydin J, Westerblad H & Lännergren J. (2003). Mitochondrial and myoplasmic $[Ca^{2+}]$ in single fibres from mouse limb muscles during repeated tetanic contractions. *J Physiol* **551**, 179-190.
- Bruton JD, Aydin J, Yamada T, Shabalina IG, Ivarsson N, Zhang SJ, Wada M, Tavi P, Nedergaard J, Katz A & Westerblad H. (2010). Increased fatigue resistance linked to Ca^{2+} -stimulated mitochondrial biogenesis in muscle fibres of cold-acclimated mice. *J Physiol* **588**, 4275-4288.
- Bruton JD, Lännergren J & Westerblad H. (1998). Effects of CO_2 -induced acidification on the fatigue resistance of single mouse muscle fibers at 28 degrees C. *J Appl Physiol* (1985) **85**, 478-483.
- Bruton JD, Place N, Yamada T, Silva JP, Andrade FH, Dahlstedt AJ, Zhang SJ, Katz A, Larsson NG & Westerblad H. (2008). Reactive oxygen species and fatigue-induced prolonged low-frequency force depression in skeletal muscle fibres of rats, mice and SOD2 overexpressing mice. *J Physiol* **586**, 175-184.
- Callahan LA, Nethery D, Stofan D, DiMarco A & Supinski G. (2001). Free radical-induced contractile protein dysfunction in endotoxin-induced sepsis. *Am J Respir Cell Mol Biol* **24**, 210-217.
- Capiati DA, Tellez-Inon MT & Boland RL. (1999). Participation of protein kinase C alpha in 1,25-dihydroxy-vitamin D_3 regulation of chick myoblast proliferation and differentiation. *Mol Cell Endocrinol* **153**, 39-45.
- Cheng AJ, Bruton JD, Lanner JT & Westerblad H. (2015). Antioxidant treatments do not improve force recovery after fatiguing stimulation of mouse skeletal muscle fibres. *J Physiol* **593**, 457-472.
- Cheng AJ, Hwee DT, Kim LH, Durham N, Yang HT, Hinken AC, Kennedy AR, Terjung RL, Jasper JR, Malik FI & Westerblad H. (2019). Fast skeletal muscle troponin activator CK-2066260 increases fatigue resistance by reducing the energetic cost of muscle contraction. *J Physiol* **597**, 4615-4625.
- Cheng AJ & Westerblad H. (2017). Mechanical isolation, and measurement of force and myoplasmic free $[Ca^{2+}]$ in fully intact single skeletal muscle fibers. *Nat Protoc* **12**, 1763-1776.
- Cheng AJ, Willis SJ, Zinner C, Chaillou T, Ivarsson N, Ørtenblad N, Lanner JT, Holmberg HC & Westerblad H. (2017). Post-exercise recovery of contractile function and endurance in humans and mice is accelerated by heating and slowed by cooling skeletal muscle. *J Physiol* **595**, 7413-7426.
- Cheng CS, Davis BN, Madden L, Bursac N & Truskey GA. (2014). Physiology and metabolism of tissue-engineered skeletal muscle. *Exp Biol Med (Maywood)* **239**, 1203-1214.

- Chin ER & Allen DG. (1997). Effects of reduced muscle glycogen concentration on force, Ca^{2+} release and contractile protein function in intact mouse skeletal muscle. *J Physiol* **498** (Pt 1), 17-29.
- Chung CJ, Wu C, Jones M, Kato TS, Dam TT, Givens RC, Templeton DL, Maurer MS, Naka Y, Takayama H, Mancini DM & Schulze PC. (2014). Reduced handgrip strength as a marker of frailty predicts clinical outcomes in patients with heart failure undergoing ventricular assist device placement. *J Card Fail* **20**, 310-315.
- Clapham DE. (2007). Calcium signaling. *Cell* **131**, 1047-1058.
- Claus RA, Bunck AC, Bockmeyer CL, Brunkhorst FM, Losche W, Kinscherf R & Deigner HP. (2005). Role of increased sphingomyelinase activity in apoptosis and organ failure of patients with severe sepsis. *FASEB J* **19**, 1719-1721.
- Conraads VM, Bosmans JM, Schuerwegh AJ, De Clerck LS, Bridts CH, Wuyts FL, Stevens WJ & Vrints CJ. (2003). Association of lipoproteins with cytokines and cytokine receptors in heart failure patients. Differences between ischaemic versus idiopathic cardiomyopathy. *Eur Heart J* **24**, 2221-2226.
- Cooke R, Franks K, Luciani GB & Pate E. (1988). The inhibition of rabbit skeletal muscle contraction by hydrogen ions and phosphate. *J Physiol* **395**, 77-97.
- Costa EM, Blau HM & Feldman D. (1986). 1,25-dihydroxyvitamin D₃ receptors and hormonal responses in cloned human skeletal muscle cells. *Endocrinology* **119**, 2214-2220.
- Cowart LA. (2010). A novel role for sphingolipid metabolism in oxidant-mediated skeletal muscle fatigue. Focus on "Sphingomyelinase stimulates oxidant signaling to weaken skeletal muscle and promote fatigue". *Am J Physiol Cell Physiol* **299**, C549-551.
- Cseri J, Szappanos H, Szigeti GP, Csernatony Z, Kovacs L & Csernoch L. (2002). A purinergic signal transduction pathway in mammalian skeletal muscle cells in culture. *Pflugers Arch* **443**, 731-738.
- Dahlstedt AJ, Katz A & Westerblad H. (2001). Role of myoplasmic phosphate in contractile function of skeletal muscle: studies on creatine kinase-deficient mice. *J Physiol* **533**, 379-388.
- Dahlstedt AJ, Katz A, Wieringa B & Westerblad H. (2000). Is creatine kinase responsible for fatigue? Studies of isolated skeletal muscle deficient in creatine kinase. *FASEB J* **14**, 982-990.
- Dahlstedt AJ & Westerblad H. (2001). Inhibition of creatine kinase reduces the rate of fatigue-induced decrease in tetanic $[\text{Ca}^{2+}]_i$ in mouse skeletal muscle. *J Physiol* **533**, 639-649.
- Dawson MJ, Gadian DG & Wilkie DR. (1978). Muscular fatigue investigated by phosphorus nuclear magnetic resonance. *Nature* **274**, 861-866.
- De Larichaudy J, Zufferli A, Serra F, Isidori AM, Naro F, Dessalle K, Desgeorges M, Piraud M, Cheillan D, Vidal H, Lefai E & Nemoz G. (2012). TNF-alpha- and tumor-induced skeletal muscle atrophy involves sphingolipid metabolism. *Skelet Muscle* **2**, 2.

- Deswal A, Petersen NJ, Feldman AM, Young JB, White BG & Mann DL. (2001). Cytokines and cytokine receptors in advanced heart failure: an analysis of the cytokine database from the Vesnarinone trial (VEST). *Circulation* **103**, 2055-2059.
- DiFranco M, Quinonez M, Shieh P, Fonarow GC, Cruz D, Deng MC, Vergara JL & Middlekauff HR. (2014). Action potential-evoked calcium release is impaired in single skeletal muscle fibers from heart failure patients. *PLoS One* **9**, e109309.
- Doehner W, Bunck AC, Rauchhaus M, von Haehling S, Brunkhorst FM, Ciccoira M, Tschope C, Ponikowski P, Claus RA & Anker SD. (2007). Secretory sphingomyelinase is upregulated in chronic heart failure: a second messenger system of immune activation relates to body composition, muscular functional capacity, and peripheral blood flow. *Eur Heart J* **28**, 821-828.
- Dolmetsch RE, Lewis RS, Goodnow CC & Healy JI. (1997). Differential activation of transcription factors induced by Ca^{2+} response amplitude and duration. *Nature* **386**, 855-858.
- Domingues-Faria C, Chanut A, Salles J, Berry A, Giraudet C, Patrac V, Denis P, Bouton K, Goncalves-Mendes N, Vasson MP, Boirie Y & Walrand S. (2014). Vitamin D deficiency down-regulates Notch pathway contributing to skeletal muscle atrophy in old wistar rats. *Nutr Metab (Lond)* **11**, 47.
- Donaldson SK, Hermansen L & Bolles L. (1978). Differential, direct effects of H^+ on Ca^{2+} -activated force of skinned fibers from the soleus, cardiac and adductor magnus muscles of rabbits. *Pflugers Arch* **376**, 55-65.
- Dridi H, Kushnir A, Zalk R, Yuan Q, Melville Z & Marks AR. (2020). Intracellular calcium leak in heart failure and atrial fibrillation: a unifying mechanism and therapeutic target. *Nat Rev Cardiol*.
- Duhamel TA, Perco JG & Green HJ. (2006). Manipulation of dietary carbohydrates after prolonged effort modifies muscle sarcoplasmic reticulum responses in exercising males. *Am J Physiol Regul Integr Comp Physiol* **291**, R1100-1110.
- Dutka TL, Mollica JP, Lambole CR, Weerakkody VC, Greening DW, Posterino GS, Murphy RM & Lamb GD. (2017). S-nitrosylation and S-glutathionylation of Cys134 on troponin I have opposing competitive actions on Ca^{2+} sensitivity in rat fast-twitch muscle fibers. *Am J Physiol Cell Physiol* **312**, C316-C327.
- Ebashi S. (1991). Excitation-contraction coupling and the mechanism of muscle contraction. *Annu Rev Physiol* **53**, 1-16.
- Edwards RH, Hill DK, Jones DA & Merton PA. (1977). Fatigue of long duration in human skeletal muscle after exercise. *J Physiol* **272**, 769-778.
- Empinado HM, Deevska GM, Nikolova-Karakashian M, Yoo JK, Christou DD & Ferreira LF. (2014). Diaphragm dysfunction in heart failure is accompanied by increases in neutral sphingomyelinase activity and ceramide content. *Eur J Heart Fail* **16**, 519-525.
- Endo I, Inoue D, Mitsui T, Umaki Y, Akaike M, Yoshizawa T, Kato S & Matsumoto T. (2003). Deletion of vitamin D receptor gene in mice results in abnormal

skeletal muscle development with deregulated expression of myoregulatory transcription factors. *Endocrinology* **144**, 5138-5144.

- Fabiato A & Fabiato F. (1978). Effects of pH on the myofilaments and the sarcoplasmic reticulum of skinned cells from cardiac and skeletal muscles. *J Physiol* **276**, 233-255.
- Ferrari R, Bachetti T, Confortini R, Opasich C, Febo O, Corti A, Cassani G & Visioli O. (1995). Tumor necrosis factor soluble receptors in patients with various degrees of congestive heart failure. *Circulation* **92**, 1479-1486.
- Ferreira LF, Moylan JS, Gilliam LA, Smith JD, Nikolova-Karakashian M & Reid MB. (2010). Sphingomyelinase stimulates oxidant signaling to weaken skeletal muscle and promote fatigue. *Am J Physiol Cell Physiol* **299**, C552-560.
- Ferreira LF, Moylan JS, Stasko S, Smith JD, Campbell KS & Reid MB. (2012). Sphingomyelinase depresses force and calcium sensitivity of the contractile apparatus in mouse diaphragm muscle fibers. *J Appl Physiol (1985)* **112**, 1538-1545.
- Fitts RH. (2016). The Role of Acidosis in Fatigue: Pro Perspective. *Med Sci Sports Exerc* **48**, 2335-2338.
- Friedrich O, Reid MB, Van den Berghe G, Vanhorebeek I, Hermans G, Rich MM & Larsson L. (2015). The Sick and the Weak: Neuropathies/Myopathies in the Critically Ill. *Physiol Rev* **95**, 1025-1109.
- Fryer MW, Owen VJ, Lamb GD & Stephenson DG. (1995). Effects of creatine phosphate and P_i on Ca^{2+} movements and tension development in rat skinned skeletal muscle fibres. *J Physiol* **482** (Pt 1), 123-140.
- Gandevia SC. (2001). Spinal and supraspinal factors in human muscle fatigue. *Physiol Rev* **81**, 1725-1789.
- Garcia LA, Ferrini MG, Norris KC & Artaza JN. (2013). $1,25(OH)_2$ vitamin D_3 enhances myogenic differentiation by modulating the expression of key angiogenic growth factors and angiogenic inhibitors in C2C12 skeletal muscle cells. *J Steroid Biochem Mol Biol* **133**, 1-11.
- Garcia LA, King KK, Ferrini MG, Norris KC & Artaza JN. (2011). $1,25(OH)_2$ vitamin D_3 stimulates myogenic differentiation by inhibiting cell proliferation and modulating the expression of promyogenic growth factors and myostatin in C2C12 skeletal muscle cells. *Endocrinology* **152**, 2976-2986.
- Gejl KD, Hvid LG, Willis SJ, Andersson E, Holmberg HC, Jensen R, Frandsen U, Hansen J, Plomgaard P & Ørtenblad N. (2016). Repeated high-intensity exercise modulates Ca^{2+} sensitivity of human skeletal muscle fibers. *Scand J Med Sci Sports* **26**, 488-497.
- Girgis CM, Clifton-Bligh RJ, Hamrick MW, Holick MF & Gunton JE. (2013). The roles of vitamin D in skeletal muscle: form, function, and metabolism. *Endocr Rev* **34**, 33-83.
- Girgis CM, Clifton-Bligh RJ, Mokbel N, Cheng K & Gunton JE. (2014a). Vitamin D signaling regulates proliferation, differentiation, and myotube size in C2C12 skeletal muscle cells. *Endocrinology* **155**, 347-357.

- Girgis CM, Mokbel N, Cha KM, Houweling PJ, Abboud M, Fraser DR, Mason RS, Clifton-Bligh RJ & Gunton JE. (2014b). The vitamin D receptor (VDR) is expressed in skeletal muscle of male mice and modulates 25-hydroxyvitamin D (25OHD) uptake in myofibers. *Endocrinology* **155**, 3227-3237.
- Gomes MD, Lecker SH, Jagoe RT, Navon A & Goldberg AL. (2001). Atrogin-1, a muscle-specific F-box protein highly expressed during muscle atrophy. *Proc Natl Acad Sci U S A* **98**, 14440-14445.
- Gopinath SD, Webb AE, Brunet A & Rando TA. (2014). FOXO3 promotes quiescence in adult muscle stem cells during the process of self-renewal. *Stem Cell Reports* **2**, 414-426.
- Gordon AM, Homsher E & Regnier M. (2000). Regulation of contraction in striated muscle. *Physiol Rev* **80**, 853-924.
- Grimby L & Hannerz J. (1977). Firing rate and recruitment order of toe extensor motor units in different modes of voluntary contraction. *J Physiol* **264**, 865-879.
- Grynkiewicz G, Poenie M & Tsien RY. (1985). A new generation of Ca^{2+} indicators with greatly improved fluorescence properties. *J Biol Chem* **260**, 3440-3450.
- Gundersen K. (2011). Excitation-transcription coupling in skeletal muscle: the molecular pathways of exercise. *Biol Rev Camb Philos Soc* **86**, 564-600.
- Hager G, Formanek M, Gedlicka C, Thurnher D, Knerer B & Kornfehl J. (2001). $1,25(\text{OH})_2$ vitamin D_3 induces elevated expression of the cell cycle-regulating genes P21 and P27 in squamous carcinoma cell lines of the head and neck. *Acta Otolaryngol* **121**, 103-109.
- Hanaoka BY, Ormseth MJ, Michael Stein C, Banerjee D, Nikolova-Karakashian M & Crofford LJ. (2018). Secretory sphingomyelinase (S-SMase) activity is elevated in patients with rheumatoid arthritis. *Clin Rheumatol* **37**, 1395-1399.
- Harber MP, Konopka AR, Douglass MD, Minchev K, Kaminsky LA, Trappe TA & Trappe S. (2009). Aerobic exercise training improves whole muscle and single myofiber size and function in older women. *Am J Physiol Regul Integr Comp Physiol* **297**, R1452-1459.
- Harrington D, Anker SD, Chua TP, Webb-Peploe KM, Ponikowski PP, Poole-Wilson PA & Coats AJ. (1997). Skeletal muscle function and its relation to exercise tolerance in chronic heart failure. *J Am Coll Cardiol* **30**, 1758-1764.
- Helliwell PS & Jackson S. (1994). Relationship between weakness and muscle wasting in rheumatoid arthritis. *Ann Rheum Dis* **53**, 726-728.
- Herrmann-Frank A, Richter M, Sarkozi S, Mohr U & Lehmann-Horn F. (1996). 4-Chloro-m-cresol, a potent and specific activator of the skeletal muscle ryanodine receptor. *Biochim Biophys Acta* **1289**, 31-40.
- Hinits Y & Hughes SM. (2007). Mef2s are required for thick filament formation in nascent muscle fibres. *Development* **134**, 2511-2519.
- Hollis BW & Wagner CL. (2005). Normal serum vitamin D levels. *N Engl J Med* **352**, 515-516; author reply 515-516.

- Huang da W, Sherman BT & Lempicki RA. (2009a). Bioinformatics enrichment tools: paths toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Res* **37**, 1-13.
- Huang da W, Sherman BT & Lempicki RA. (2009b). Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc* **4**, 44-57.
- Hulsmann M, Quittan M, Berger R, Crevenna R, Springer C, Nuhr M, Mortl D, Moser P & Pacher R. (2004). Muscle strength as a predictor of long-term survival in severe congestive heart failure. *Eur J Heart Fail* **6**, 101-107.
- Hvid LG, Gejl K, Bech RD, Nygaard T, Jensen K, Frandsen U & Ørtenblad N. (2013). Transient impairments in single muscle fibre contractile function after prolonged cycling in elite endurance athletes. *Acta Physiol (Oxf)* **208**, 265-273.
- Hvid LG, Ørtenblad N, Aagaard P, Kjaer M & Suetta C. (2011). Effects of ageing on single muscle fibre contractile function following short-term immobilisation. *J Physiol* **589**, 4745-4757.
- Hyde R, Hajdуч E, Powell DJ, Taylor PM & Hundal HS. (2005). Ceramide down-regulates System A amino acid transport and protein synthesis in rat skeletal muscle cells. *FASEB J* **19**, 461-463.
- Irazaqui AP, Boland RL & Buitrago CG. (2014). Actions of 1,25(OH)₂-vitamin D₃ on the cellular cycle depend on VDR and p38 MAPK in skeletal muscle cells. *J Mol Endocrinol* **53**, 331-343.
- Jenkins RW, Canals D, Idkowiak-Baldys J, Simbari F, Roddy P, Perry DM, Kitatani K, Luberto C & Hannun YA. (2010). Regulated secretion of acid sphingomyelinase: implications for selectivity of ceramide formation. *The J Biol Chem* **285**, 35706-35718.
- Jensen SS, Madsen MW, Lukas J, Binderup L & Bartek J. (2001). Inhibitory effects of 1 α ,25-dihydroxyvitamin D₃ on the G₁-S phase-controlling machinery. *Mol Endocrinol* **15**, 1370-1380.
- Khodabukus A, Madden L, Prabhu NK, Koves TR, Jackam CP, Muoio DM & Bursac N. (2019). Electrical stimulation increases hypertrophy and metabolic flux in tissue-engineered human skeletal muscle. *Biomaterials* **198**, 259-269.
- Kreutz M & Andreessen R. (1990). Induction of human monocyte to macrophage maturation in vitro by 1,25-dihydroxyvitamin D₃. *Blood* **76**, 2457-2461.
- Kreutz M, Andreessen R, Krause SW, Szabo A, Ritz E & Reichel H. (1993). 1,25-dihydroxyvitamin D₃ production and vitamin D₃ receptor expression are developmentally regulated during differentiation of human monocytes into macrophages. *Blood* **82**, 1300-1307.
- Kuang S, Kuroda K, Le Grand F & Rudnicki MA. (2007). Asymmetric self-renewal and commitment of satellite stem cells in muscle. *Cell* **129**, 999-1010.
- Kushnir A, Todd JJ, Witherspoon JW, Yuan Q, Reiken S, Lin H, Munce RH, Wajsborg B, Melville Z, Clarke OB, Wedderburn-Pugh K, Wronska A, Razaqyar MS, Chrismer IC, Shelton MO, Mankodi A, Grunseich C, Tarnopolsky MA, Tanji

- K, Hirano M, Riazi S, Kraeva N, Voermans NC, Gruber A, Allen C, Meilleur KG & Marks AR. (2020). Intracellular calcium leak as a therapeutic target for RYR1-related myopathies. *Acta Neuropathol* **139**, 1089-1104.
- Lamb GD. (2002). Excitation-contraction coupling and fatigue mechanisms in skeletal muscle: studies with mechanically skinned fibres. *J Muscle Res Cell Motil* **23**, 81-91.
- Lamb GD & Westerblad H. (2011). Acute effects of reactive oxygen and nitrogen species on the contractile function of skeletal muscle. *J Physiol* **589**, 2119-2127.
- Lambernd S, Taube A, Schober A, Platzbecker B, Gorgens SW, Schlich R, Jeruschke K, Weiss J, Eckardt K & Eckel J. (2012). Contractile activity of human skeletal muscle cells prevents insulin resistance by inhibiting pro-inflammatory signalling pathways. *Diabetologia* **55**, 1128-1139.
- Lambole CR, Rouffet DM, Dutka TL, McKenna MJ & Lamb GD. (2020). Effects of high-intensity intermittent exercise on the contractile properties of human type I and type II skeletal muscle fibers. *J Appl Physiol (1985)* **128**, 1207-1216.
- Lambole CR, Wyckelsma VL, Dutka TL, McKenna MJ, Murphy RM & Lamb GD. (2015). Contractile properties and sarcoplasmic reticulum calcium content in type I and type II skeletal muscle fibres in active aged humans. *J Physiol* **593**, 2499-2514.
- Lanner JT, Katz A, Tavi P, Sandström ME, Zhang SJ, Wretman C, James S, Fauconnier J, Lännergren J, Bruton JD & Westerblad H. (2006). The role of Ca^{2+} influx for insulin-mediated glucose uptake in skeletal muscle. *Diabetes* **55**, 2077-2083.
- Lännergren J & Westerblad H. (1987). The temperature dependence of isometric contractions of single, intact fibres dissected from a mouse foot muscle. *J Physiol* **390**, 285-293.
- Larkin LM, Van der Meulen JH, Dennis RG & Kennedy JB. (2006). Functional evaluation of nerve-skeletal muscle constructs engineered in vitro. *In Vitro Cell Dev Biol Anim* **42**, 75-82.
- Lecker SH, Jagoe RT, Gilbert A, Gomes M, Baracos V, Bailey J, Price SR, Mitch WE & Goldberg AL. (2004). Multiple types of skeletal muscle atrophy involve a common program of changes in gene expression. *FASEB J* **18**, 39-51.
- Levine B, Kalman J, Mayer L, Fillit HM & Packer M. (1990). Elevated circulating levels of tumor necrosis factor in severe chronic heart failure. *N Engl J Med* **323**, 236-241.
- Lipkin DP, Jones DA, Round JM & Poole-Wilson PA. (1988). Abnormalities of skeletal muscle in patients with chronic heart failure. *Int J Cardiol* **18**, 187-195.
- Llano-Diez M, Cheng AJ, Jonsson W, Ivarsson N, Westerblad H, Sun V, Cacciani N, Larsson L & Bruton J. (2016). Impaired Ca^{2+} release contributes to muscle weakness in a rat model of critical illness myopathy. *Crit Care* **20**, 254.
- Loehr JA, Abo-Zahrah R, Pal R & Rodney GG. (2014). Sphingomyelinase promotes oxidant production and skeletal muscle contractile dysfunction through activation of NADPH oxidase. *Front Physiol* **5**, 530.

- Luden N, Minchev K, Hayes E, Louis E, Trappe T & Trappe S. (2008). Human vastus lateralis and soleus muscles display divergent cellular contractile properties. *Am J Physiol Regul Integr Comp Physiol* **295**, R1593-1598.
- Macdonald WA & Stephenson DG. (2006). Effect of ADP on slow-twitch muscle fibres of the rat: implications for muscle fatigue. *J Physiol* **573**, 187-198.
- Madden L, Juhas M, Kraus WE, Truskey GA & Bursac N. (2015). Bioengineered human myobundles mimic clinical responses of skeletal muscle to drugs. *Elife* **4**.
- Malisoux L, Francaux M, Nielens H, Renard P, Lebacqz J & Theisen D. (2006). Calcium sensitivity of human single muscle fibers following plyometric training. *Med Sci Sports Exerc* **38**, 1901-1908.
- Marathe S, Schissel SL, Yellin MJ, Beatini N, Mintzer R, Williams KJ & Tabas I. (1998). Human vascular endothelial cells are a rich and regulatable source of secretory sphingomyelinase. Implications for early atherogenesis and ceramide-mediated cell signaling. *J Biol Chem* **273**, 4081-4088.
- Marsden CD, Meadows JC & Merton PA. (1971). Isolated single motor units in human muscle and their rate of discharge during maximal voluntary effort. *J Physiol* **217 Suppl**, 12P-13P.
- Martin NR, Passey SL, Player DJ, Khodabukus A, Ferguson RA, Sharples AP, Mudera V, Baar K & Lewis MP. (2013). Factors affecting the structure and maturation of human tissue engineered skeletal muscle. *Biomaterials* **34**, 5759-5765.
- Mauro A. (1961). Satellite cell of skeletal muscle fibers. *J Biophys Biochem Cytol* **9**, 493-495.
- McKenna CF & Fry CS. (2017). Altered satellite cell dynamics accompany skeletal muscle atrophy during chronic illness, disuse, and aging. *Curr Opin Clin Nutr Metab Care* **20**, 447-452.
- McPherron AC, Lawler AM & Lee SJ. (1997). Regulation of skeletal muscle mass in mice by a new TGF-beta superfamily member. *Nature* **387**, 83-90.
- Mebarek S, Komati H, Naro F, Zeiller C, Alvisi M, Lagarde M, Prigent AF & Nemoz G. (2007). Inhibition of de novo ceramide synthesis upregulates phospholipase D and enhances myogenic differentiation. *J Cell Sci* **120**, 407-416.
- Menconi MJ, Wei W, Yang H, Wray CJ & Hasselgren PO. (2004). Treatment of cultured myotubes with the calcium ionophore A23187 increases proteasome activity via a CaMK II-caspase-calpain-dependent mechanism. *Surgery* **136**, 135-142.
- Merton PA. (1954). Voluntary strength and fatigue. *J Physiol* **123**, 553-564.
- Middlekauff HR. (2010). Making the case for skeletal myopathy as the major limitation of exercise capacity in heart failure. *Circ Heart Fail* **3**, 537-546.
- Middlekauff HR, Vigna C, Verity MA, Fonarow GC, Horwich TB, Hamilton MA, Shieh P & Tupling AR. (2012). Abnormalities of calcium handling proteins in

skeletal muscle mirror those of the heart in humans with heart failure: a shared mechanism? *J Card Fail* **18**, 724-733.

- Millar NC & Homsher E. (1990). The effect of phosphate and calcium on force generation in glycerinated rabbit skeletal muscle fibers. A steady-state and transient kinetic study. *J Biol Chem* **265**, 20234-20240.
- Mizuno M. (1991). Human respiratory muscles: fibre morphology and capillary supply. *Eur Respir J* **4**, 587-601.
- Mizuno M & Secher NH. (1989). Histochemical characteristics of human expiratory and inspiratory intercostal muscles. *J Appl Physiol (1985)* **67**, 592-598.
- Mizuno M, Secher NH & Saltin B. (1985). Fibre types, capillary supply and enzyme activities in human intercostal muscles. *Clin Physiol* **5**, 121-135.
- Mollica JP, Dutka TL, Merry TL, Lambole CR, McConell GK, McKenna MJ, Murphy RM & Lamb GD. (2012). S-glutathionylation of troponin I (fast) increases contractile apparatus Ca^{2+} sensitivity in fast-twitch muscle fibres of rats and humans. *J Physiol* **590**, 1443-1463.
- Moylan JS, Smith JD, Wolf Horrell EM, McLean JB, Deevska GM, Bonnell MR, Nikolova-Karakashian MN & Reid MB. (2014). Neutral sphingomyelinase-3 mediates TNF-stimulated oxidant activity in skeletal muscle. *Redox Biol* **2**, 910-920.
- Murphy RM, Dutka TL & Lamb GD. (2008). Hydroxyl radical and glutathione interactions alter calcium sensitivity and maximum force of the contractile apparatus in rat skeletal muscle fibres. *J Physiol* **586**, 2203-2216.
- Need AG, O'Loughlin PD, Morris HA, Coates PS, Horowitz M & Nordin BE. (2008). Vitamin D metabolites and calcium absorption in severe vitamin D deficiency. *J Bone Miner Res* **23**, 1859-1863.
- Needleman DH, Aghdasi B, Seryshev AB, Schroepfer GJ, Jr. & Hamilton SL. (1997). Modulation of skeletal muscle Ca^{2+} -release channel activity by sphingosine. *Am J Physiol* **272**, C1465-1474.
- Nelson CR, Debold EP & Fitts RH. (2014). Phosphate and acidosis act synergistically to depress peak power in rat muscle fibers. *Am J Physiol Cell Physiol* **307**, C939-950.
- Nelson CR & Fitts RH. (2014). Effects of low cell pH and elevated inorganic phosphate on the pCa-force relationship in single muscle fibers at near-physiological temperatures. *Am J Physiol Cell Physiol* **306**, C670-678.
- Nielsen J, Cheng AJ, Ørtenblad N & Westerblad H. (2014). Subcellular distribution of glycogen and decreased tetanic Ca^{2+} in fatigued single intact mouse muscle fibres. *J Physiol* **592**, 2003-2012.
- Nielsen J, Schroder HD, Rix CG & Ørtenblad N. (2009). Distinct effects of subcellular glycogen localization on tetanic relaxation time and endurance in mechanically skinned rat skeletal muscle fibres. *J Physiol* **587**, 3679-3690.

- Nikolic N, Bakke SS, Kase ET, Rudberg I, Flo Halle I, Rustan AC, Thoresen GH & Aas V. (2012). Electrical pulse stimulation of cultured human skeletal muscle cells as an in vitro model of exercise. *PLoS One* **7**, e33203.
- Nikolova-Karakashian MN & Reid MB. (2011). Sphingolipid metabolism, oxidant signaling, and contractile function of skeletal muscle. *Antioxid Redox Signal* **15**, 2501-2517.
- Norman B, Hellsten-Westling Y, Sjödin B & Jansson E. (1994). AMP deaminase in skeletal muscle of healthy males quantitatively determined by new assay. *Acta Physiol Scand* **150**, 397-403
- Okuno H, Kishimoto KN, Hatori M & Itoi E. (2012). 1alpha,25-dihydroxyvitamin D₃ enhances fast-myosin heavy chain expression in differentiated C2C12 myoblasts. *Cell Biol Int* **36**, 441-447.
- Olsson K, Cheng AJ, Alam S, Al-Ameri M, Rullman E, Westerblad H, Lanner JT, Bruton JD & Gustafsson T. (2015). Intracellular Ca²⁺-handling differs markedly between intact human muscle fibers and myotubes. *Skelet Muscle* **5**, 26.
- Ørtenblad N, Nielsen J, Saltin B & Holmberg HC. (2011). Role of glycogen availability in sarcoplasmic reticulum Ca²⁺ kinetics in human skeletal muscle. *J Physiol* **589**, 711-725.
- Park YG, Moon JH & Kim J. (2006). A comparative study of magnetic-activated cell sorting, cytotoxicity and preplating for the purification of human myoblasts. *Yonsei Med J* **47**, 179-183.
- Pate E, Bhimani M, Franks-Skiba K & Cooke R. (1995). Reduced effect of pH on skinned rabbit psoas muscle mechanics at high temperatures: implications for fatigue. *J Physiol* **486** (Pt 3), 689-694.
- Pate E & Cooke R. (1989). Addition of phosphate to active muscle fibers probes actomyosin states within the powerstroke. *Pflugers Arch* **414**, 73-81.
- Perreault CL, Gonzalez-Serratos H, Litwin SE, Sun X, Franzini-Armstrong C & Morgan JP. (1993). Alterations in contractility and intracellular Ca²⁺ transients in isolated bundles of skeletal muscle fibers from rats with chronic heart failure. *Circ Res* **73**, 405-412.
- Peter A, Weigert C, Staiger H, Machicao F, Schick F, Machann J, Stefan N, Thamer C, Haring HU & Schleicher E. (2009). Individual stearyl-coa desaturase 1 expression modulates endoplasmic reticulum stress and inflammation in human myotubes and is associated with skeletal muscle lipid storage and insulin sensitivity in vivo. *Diabetes* **58**, 1757-1765.
- Philipp S, Puchert M, Adam-Klages S, Tchikov V, Winoto-Morbach S, Mathieu S, Deerberg A, Kolker L, Marchesini N, Kabelitz D, Hannun YA, Schutze S & Adam D. (2010). The Polycomb group protein EED couples TNF receptor 1 to neutral sphingomyelinase. *Proc Natl Acad Sci U S A* **107**, 1112-1117.
- Phillips BE, Hill DS & Atherton PJ. (2012). Regulation of muscle protein synthesis in humans. *Curr Opin Clin Nutr Metab Care* **15**, 58-63.

- Pike JW. (2014). Expression of the vitamin D receptor in skeletal muscle: are we there yet? *Endocrinology* **155**, 3214-3218.
- Place N, Ivarsson N, Venckunas T, Neyroud D, Brazaitis M, Cheng AJ, Ochala J, Kamandulis S, Girard S, Volungevicius G, Pauzas H, Mekideche A, Kayser B, Martinez-Redondo V, Ruas JL, Bruton J, Truffert A, Lanner JT, Skurvydas A & Westerblad H. (2015). Ryanodine receptor fragmentation and sarcoplasmic reticulum Ca^{2+} leak after one session of high-intensity interval exercise. *Proc Natl Acad Sci U S A* **112**, 15492-15497.
- Place N, Yamada T, Bruton JD & Westerblad H. (2010). Muscle fatigue: from observations in humans to underlying mechanisms studied in intact single muscle fibres. *Eur J Appl Physiol* **110**, 1-15.
- Pleasure D, Wyszynski B, Sumner A, Schotland D, Feldman B, Nugent N, Hitz K & Goodman DB. (1979). Skeletal muscle calcium metabolism and contractile force in vitamin D-deficient chicks. *J Clin Invest* **64**, 1157-1167.
- Pojednic RM & Ceglia L. (2014). The emerging biomolecular role of vitamin D in skeletal muscle. *Exerc Sport Sci Rev* **42**, 76-81.
- Pojednic RM, Ceglia L, Olsson K, Gustafsson T, Lichtenstein AH, Dawson-Hughes B & Fielding RA. (2015). Effects of 1,25-dihydroxyvitamin D_3 and vitamin D_3 on the expression of the vitamin d receptor in human skeletal muscle cells. *Calcif Tissue Int* **96**, 256-263.
- Posterino GS, Lamb GD & Stephenson DG. (2000). Twitch and tetanic force responses and longitudinal propagation of action potentials in skinned skeletal muscle fibres of the rat. *J Physiol* **527 Pt 1**, 131-137.
- Potthoff MJ, Arnold MA, McAnally J, Richardson JA, Bassel-Duby R & Olson EN. (2007). Regulation of skeletal muscle sarcomere integrity and postnatal muscle function by Mef2c. *Mol Cell Biol* **27**, 8143-8151.
- Powers SK, Lynch GS, Murphy KT, Reid MB & Zijdwind I. (2016). Disease-Induced Skeletal Muscle Atrophy and Fatigue. *Med Sci Sports Exerc* **48**, 2307-2319.
- Rao L, Qian Y, Khodabukus A, Ribar T & Bursac N. (2018). Engineering human pluripotent stem cells into functional skeletal muscle tissue. *Nat Commun* **9**, 126
- Raschke S, Eckardt K, Bjørklund Holven K, Jensen J & Eckel J. (2013a). Identification and validation of novel contraction-regulated myokines released from primary human skeletal muscle cells. *PLoS One* **8**, e62008.
- Raschke S, Elsen M, Gassenhuber H, Sommerfeld M, Schwahn U, Brockmann B, Jung R, Wisloff U, Tjonna AE, Raastad T, Hallen J, Norheim F, Drevon CA, Romacho T, Eckardt K & Eckel J. (2013b). Evidence against a beneficial effect of irisin in humans. *PLoS One* **8**, e73680.
- Rauchhaus M, Doehner W, Francis DP, Davos C, Kemp M, Liebenthal C, Niebauer J, Hooper J, Volk HD, Coats AJ & Anker SD. (2000). Plasma cytokine parameters and mortality in patients with chronic heart failure. *Circulation* **102**, 3060-3067.

- Rebbbeck RT, Karunasekara Y, Board PG, Beard NA, Casarotto MG & Dulhunty AF. (2014). Skeletal muscle excitation-contraction coupling: who are the dancing partners? *Int J Biochem Cell Biol* **48**, 28-38.
- Reid MB, Lännergren J & Westerblad H. (2002). Respiratory and limb muscle weakness induced by tumor necrosis factor-alpha: involvement of muscle myofilaments. *Am J Resp Crit Care Med* **166**, 479-484.
- Reid MB & Moylan JS. (2011). Beyond atrophy: redox mechanisms of muscle dysfunction in chronic inflammatory disease. *J Physiol* **589**, 2171-2179.
- Reiken S, Lacampagne A, Zhou H, Kherani A, Lehnart SE, Ward C, Huang F, Gaburjakova M, Gaburjakova J, Rosemblyt N, Warren MS, He KL, Yi GH, Wang J, Burkhoff D, Vassort G & Marks AR. (2003). PKA phosphorylation activates the calcium release channel (ryanodine receptor) in skeletal muscle: defective regulation in heart failure. *J Cell Biol* **160**, 919-928.
- Relaix F & Zammit PS. (2012). Satellite cells are essential for skeletal muscle regeneration: the cell on the edge returns centre stage. *Development* **139**, 2845-2856.
- Rodman JS & Baker T. (1978). Changes in the kinetics of muscle contraction in vitamin D-depleted rats. *Kidney Int* **13**, 189-193.
- Rokach O, Ullrich ND, Rausch M, Mouly V, Zhou H, Muntoni F, Zorzato F & Treves S. (2013). Establishment of a human skeletal muscle-derived cell line: biochemical, cellular and electrophysiological characterization. *Biochem J* **455**, 169-177.
- Rousseau E, Ladine J, Liu QY & Meissner G. (1988). Activation of the Ca^{2+} release channel of skeletal muscle sarcoplasmic reticulum by caffeine and related compounds. *Arch Biochem Biophys* **267**, 75-86.
- Ruff RL & Whittlesey D. (1991). Ca-, Sr-tension relationships and contraction velocities of human muscle fibers. *Muscle Nerve* **14**, 1219-1226.
- Rullman E, Andersson DC, Melin M, Reiken S, Mancini DM, Marks AR, Lund LH & Gustafsson T. (2013). Modifications of skeletal muscle ryanodine receptor type 1 and exercise intolerance in heart failure. *J Heart Lung Transplant* **32**, 925-929.
- Rullman E, Melin M, Mandic M, Gonon A, Fernandez-Gonzalo R & Gustafsson T. (2020). Circulatory factors associated with function and prognosis in patients with severe heart failure. *Clin Res Cardiol* **109**, 655-672.
- Sabbadini RA, Betto R, Teresi A, Fachechi-Cassano G & Salviati G. (1992). The effects of sphingosine on sarcoplasmic reticulum membrane calcium release. *J Biol Chem* **267**, 15475-15484.
- Sahlin K, Harris RC & Hultman E. (1975). Creatine kinase equilibrium and lactate content compared with muscle pH in tissue samples obtained after isometric exercise. *Biochem J* **152**, 173-180.
- Sahlin K, Harris RC, Nyling B & Hultman E. (1976). Lactate content and pH in muscle obtained after dynamic exercise. *Pflugers Arch* **367**, 143-149.

- Sala M, Hollinger KR, Thomas AG, Dash RP, Tallon C, Veeravalli V, Lovell L, Kogler M, Hrebabecky H, Prochazkova E, Nesuta O, Donoghue A, Lam J, Rais R, Rojas C, Slusher BS & Nencka R. (2020). Novel Human Neutral Sphingomyelinase 2 Inhibitors as Potential Therapeutics for Alzheimer's Disease. *J Med Chem* **63**, 6028-6056.
- Sapega AA, Sokolow DP, Graham TJ & Chance B. (1987). Phosphorus nuclear magnetic resonance: a non-invasive technique for the study of muscle bioenergetics during exercise. *Med Sci Sports Exerc* **19**, 410-420.
- Sarabia V, Lam L, Burdett E, Leiter LA & Klip A. (1992). Glucose transport in human skeletal muscle cells in culture. Stimulation by insulin and metformin. *J Clin Invest* **90**, 1386-1395.
- Schantz P, Randall-Fox E, Hutchison W, Tyden A & Åstrand PO. (1983). Muscle fibre type distribution, muscle cross-sectional area and maximal voluntary strength in humans. *Acta Physiol Scand* **117**, 219-226.
- Scheler M, Irmeler M, Lehr S, Hartwig S, Staiger H, Al-Hasani H, Beckers J, de Angelis MH, Haring HU & Weigert C. (2013). Cytokine response of primary human myotubes in an in vitro exercise model. *Am J Physiol Cell Physiol* **305**, C877-886.
- Shen B, Arese M, Gualandris A & Rifkin DB. (1998). Intracellular association of FGF-2 with the ribosomal protein L6/TAXREB107. *Biochem Biophys Res Commun* **252**, 524-528.
- Simpson RU, Thomas GA & Arnold AJ. (1985). Identification of 1,25-dihydroxyvitamin D₃ receptors and activities in muscle. *J Biol Chem* **260**, 8882-8891.
- Sjaland C, Lunde PK, Swift F, Munkvik M, Ericsson M, Lunde M, Boye S, Christensen G, Ellingsen O, Sejersted OM & Andersson KB. (2011). Slowed relaxation and preserved maximal force in soleus muscles of mice with targeted disruption of the Serca2 gene in skeletal muscle. *J Physiol* **589**, 6139-6155.
- Skurvydas A, Mamkus G, Kamandulis S, Dudonienė V, Valancienė D & Westerblad H. (2016). Mechanisms of force depression caused by different types of physical exercise studied by direct electrical stimulation of human quadriceps muscle. *Eur J Appl Physiol* **116**, 2215-2224.
- Sluzalska KD, Slawski J, Sochacka M, Lampart A, Otlewski J & Zakrzewska M. (2020). Intracellular partners of fibroblast growth factors 1 and 2 - implications for functions. *Cytokine Growth Factor Rev*.
- Smith AS, Long CJ, Pirozzi K, Najjar S, McAleer C, Vandeburgh HH & Hickman JJ. (2014). A multiplexed chip-based assay system for investigating the functional development of human skeletal myotubes in vitro. *J Biotechnol* **185**, 15-18.
- Smith IJ & Dodd SL. (2007). Calpain activation causes a proteasome-dependent increase in protein degradation and inhibits the Akt signalling pathway in rat diaphragm muscle. *Exp Physiol* **92**, 561-573.
- Sokka T, Hakkinen A, Kautiainen H, Maillefert JF, Toloza S, Mork Hansen T, Calvo-Alen J, Oding R, Liveborn M, Huisman M, Alten R, Pohl C, Cutolo M, Immonen K, Woolf A, Murphy E, Sheehy C, Quirke E, Celik S, Yazici Y,

- Tlustochowicz W, Kapolka D, Skakic V, Rojkovich B, Muller R, Stropuviene S, Andersone D, Drosos AA, Lazovskis J, Pincus T & Group Q-R. (2008). Physical inactivity in patients with rheumatoid arthritis: data from twenty-one countries in a cross-sectional, international study. *Arth Rheum* **59**, 42-50.
- Srikuea R, Zhang X, Park-Sarge OK & Esser KA. (2012). VDR and CYP27B1 are expressed in C2C12 cells and regenerating skeletal muscle: potential role in suppression of myoblast proliferation. *Am J Physiol Cell Physiol* **303**, C396-405.
- Stasko SA, Hardin BJ, Smith JD, Moylan JS & Reid MB. (2013). TNF signals via neuronal-type nitric oxide synthase and reactive oxygen species to depress specific force of skeletal muscle. *J Appl Physiol (1985)* **114**, 1629-1636.
- Steinz MM, Persson M, Aresh B, Olsson K, Cheng AJ, Ahlstrand E, Lilja M, Lundberg TR, Rullman E, Angeby Moller K, Sandor K, Ajeganova S, Yamada T, Beard N, Karlsson BC, Tavi P, Kenne E, Svensson CI, Rassier DE, Karlsson R, Friedman R, Gustafsson T & Lanner JT. (2019). Oxidative hotspots on actin promote skeletal muscle weakness in rheumatoid arthritis. *JCI Insight* **5**.
- Strle K, Broussard SR, McCusker RH, Shen WH, Johnson RW, Freund GG, Dantzer R & Kelley KW. (2004). Proinflammatory cytokine impairment of insulin-like growth factor I-induced protein synthesis in skeletal muscle myoblasts requires ceramide. *Endocrinology* **145**, 4592-4602.
- Sundberg CW, Hunter SK, Trappe SW, Smith CS & Fitts RH. (2018). Effects of elevated H⁺ and P_i on the contractile mechanics of skeletal muscle fibres from young and old men: implications for muscle fatigue in humans. *J Physiol* **596**, 3993-4015.
- Supinski GS, Alimov AP, Wang L, Song XH & Callahan LA. (2015). Neutral sphingomyelinase 2 is required for cytokine-induced skeletal muscle calpain activation. *Am J Physiol Lung Cell Mol Physiol* **309**, L614-624.
- Sweeney HL, Bowman BF & Stull JT. (1993). Myosin light chain phosphorylation in vertebrate striated muscle: regulation and function. *Am J Physiol* **264**, C1085-1095.
- Tanabe T, Beam KG, Adams BA, Niidome T & Numa S. (1990). Regions of the skeletal muscle dihydropyridine receptor critical for excitation-contraction coupling. *Nature* **346**, 567-569.
- Tanaka H, Furuya T, Kameda N, Kobayashi T & Mizusawa H. (2000). Triad proteins and intracellular Ca²⁺ transients during development of human skeletal muscle cells in aneural and innervated cultures. *J Muscl Res Cell Motil* **21**, 507-526.
- Tanaka K, Kanazawa I, Yamaguchi T, Yano S, Kaji H & Sugimoto T. (2014a). Active vitamin D possesses beneficial effects on the interaction between muscle and bone. *Biochem Biophys Res Commun* **450**, 482-487.
- Tanaka M, Kishimoto KN, Okuno H, Saito H & Itoi E. (2014b). Vitamin D receptor gene silencing effects on differentiation of myogenic cell lines. *Muscle Nerve* **49**, 700-708.

- Tavi P & Westerblad H. (2011). The role of in vivo Ca^{2+} signals acting on Ca^{2+} -calmodulin-dependent proteins for skeletal muscle plasticity. *J Physiol* **589**, 5021-5031.
- Taylor JL, Todd G & Gandevia SC. (2006). Evidence for a supraspinal contribution to human muscle fatigue. *Clin Exp Pharm Physiol* **33**, 400-405.
- Testa M, Yeh M, Lee P, Fanelli R, Loperfido F, Berman JW & LeJemtel TH. (1996). Circulating levels of cytokines and their endogenous modulators in patients with mild to severe congestive heart failure due to coronary artery disease or hypertension. *J Am Coll Cardiol* **28**, 964-971.
- Torre-Amione G, Kapadia S, Benedict C, Oral H, Young JB & Mann DL. (1996). Proinflammatory cytokine levels in patients with depressed left ventricular ejection fraction: a report from the Studies of Left Ventricular Dysfunction (SOLVD). *J Am Coll Cardiol* **27**, 1201-1206.
- Toth MJ, Ades PA, Tischler MD, Tracy RP & LeWinter MM. (2006). Immune activation is associated with reduced skeletal muscle mass and physical function in chronic heart failure. *Int J Card* **109**, 179-187.
- Toth MJ, Miller MS, Ward KA & Ades PA. (2012). Skeletal muscle mitochondrial density, gene expression, and enzyme activities in human heart failure: minimal effects of the disease and resistance training. *J Appl Physiol* (1985) **112**, 1864-1874.
- Toth MJ, Shaw AO, Miller MS, VanBuren P, LeWinter MM, Maughan DW & Ades PA. (2010). Reduced knee extensor function in heart failure is not explained by inactivity. *Int J Card* **143**, 276-282.
- Tracey KJ, Lowry SF, Beutler B, Cerami A, Albert JD & Shires GT. (1986). Cachectin/tumor necrosis factor mediates changes of skeletal muscle plasma membrane potential. *J Exp Med* **164**, 1368-1373.
- Ullrich ND, Fischer D, Kornblum C, Walter MC, Niggli E, Zorzato F & Treves S. (2011). Alterations of excitation-contraction coupling and excitation coupled Ca^{2+} entry in human myotubes carrying CAV3 mutations linked to rippling muscle. *Hum Mutat* **32**, 309-317.
- Wang Y, Becklund BR & DeLuca HF. (2010). Identification of a highly specific and versatile vitamin D receptor antibody. *Arch Biochem Biophys* **494**, 166-177.
- Wang Y & DeLuca HF. (2011). Is the vitamin d receptor found in muscle? *Endocrinology* **152**, 354-363.
- Ward CW, Reiken S, Marks AR, Marty I, Vassort G & Lacampagne A. (2003). Defects in ryanodine receptor calcium release in skeletal muscle from post-myocardial infarct rats. *FASEB J* **17**, 1517-1519.
- Watanabe D, Aibara C & Wada M. (2019). Treatment with EUK-134 improves sarcoplasmic reticulum Ca^{2+} release but not myofibrillar Ca^{2+} sensitivity after fatiguing contraction of rat fast-twitch muscle. *Am J Physiol Reg Integr Comp Physiol* **316**, R543-R551.
- Watanabe D, Kanzaki K, Kuratani M, Matsunaga S, Yanaka N & Wada M. (2015). Contribution of impaired myofibril and ryanodine receptor function to

- prolonged low-frequency force depression after in situ stimulation in rat skeletal muscle. *J Muscle Res Cell Motil* **36**, 275-286.
- Watanabe D & Wada M. (2016). Predominant cause of prolonged low-frequency force depression changes during recovery after in situ fatiguing stimulation of rat fast-twitch muscle. *Am J Physiol Reg Integr Comp Physiol* **311**, R919-R929.
- Webster RJ, Sabbadini RA, Dettbarn CA & Paolini PJ. (1994). Sphingosine effects on the contractile behavior of skinned cardiac myocytes. *J Mol Cell Cardiol* **26**, 1273-1290.
- Welinder C & Ekblad L. (2011). Coomassie staining as loading control in Western blot analysis. *J Proteome Res* **10**, 1416-1419.
- Westerblad H & Allen DG. (1991). Changes of myoplasmic calcium concentration during fatigue in single mouse muscle fibers. *J Gen Physiol* **98**, 615-635.
- Westerblad H & Allen DG. (1993a). The contribution of $[Ca^{2+}]_i$ to the slowing of relaxation in fatigued single fibres from mouse skeletal muscle. *J Physiol* **468**, 729-740.
- Westerblad H & Allen DG. (1993b). The influence of intracellular pH on contraction, relaxation and $[Ca^{2+}]_i$ in intact single fibres from mouse muscle. *J Physiol* **466**, 611-628.
- Westerblad H & Allen DG. (1996). The effects of intracellular injections of phosphate on intracellular calcium and force in single fibres of mouse skeletal muscle. *Pflugers Arch* **431**, 964-970.
- Westerblad H, Bruton JD & Lännergren J. (1997). The effect of intracellular pH on contractile function of intact, single fibres of mouse muscle declines with increasing temperature. *J Physiol* **500** (Pt 1), 193-204.
- Westerblad H, Duty S & Allen DG. (1993). Intracellular calcium concentration during low-frequency fatigue in isolated single fibers of mouse skeletal muscle. *J Appl Physiol (1985)* **75**, 382-388.
- Wilkinson DJ, Piasecki M & Atherton PJ. (2018). The age-related loss of skeletal muscle mass and function: Measurement and physiology of muscle fibre atrophy and muscle fibre loss in humans. *Ageing Res Rev* **47**, 123-132.
- Wong ML, Xie B, Beatini N, Phu P, Marathe S, Johns A, Gold PW, Hirsch E, Williams KJ, Licinio J & Tabas I. (2000). Acute systemic inflammation up-regulates secretory sphingomyelinase in vivo: a possible link between inflammatory cytokines and atherogenesis. *Proc Natl Acad Sci U S A* **97**, 8681-8686.
- Yahiaoui L, Gvozdic D, Danialou G, Mack M & Petrof BJ. (2008). CC family chemokines directly regulate myoblast responses to skeletal muscle injury. *J Physiol* **586**, 3991-4004.
- Yamada T, Abe M, Lee J, Tatebayashi D, Himori K, Kanzaki K, Wada M, Bruton JD, Westerblad H & Lanner JT. (2015a). Muscle dysfunction associated with adjuvant-induced arthritis is prevented by antioxidant treatment. *Skelet Muscle* **5**, 20.

- Yamada T, Fedotovskaya O, Cheng AJ, Cornachione AS, Minozzo FC, Aulin C, Friden C, Turesson C, Andersson DC, Glenmark B, Lundberg IE, Rassier DE, Westerblad H & Lanner JT. (2015b). Nitrosative modifications of the Ca^{2+} release complex and actin underlie arthritis-induced muscle weakness. *Ann Rheum Dis* **74**, 1907-1914.
- Yamada T, Place N, Kosterina N, Ostberg T, Zhang SJ, Grundtman C, Erlandsson-Harris H, Lundberg IE, Glenmark B, Bruton JD & Westerblad H. (2009). Impaired myofibrillar function in the soleus muscle of mice with collagen-induced arthritis. *Arthritis Rheum* **60**, 3280-3289.
- Yamada T, Steinz MM, Kenne E & Lanner JT. (2017). Muscle Weakness in Rheumatoid Arthritis: The Role of Ca^{2+} and Free Radical Signaling. *EBioMedicine* **23**, 12-19.
- Yokoyama T, Vaca L, Rossen RD, Durante W, Hazarika P & Mann DL. (1993). Cellular basis for the negative inotropic effects of tumor necrosis factor- α in the adult mammalian heart. *J Clin Invest* **92**, 2303-2312.